

## Comparison of effectiveness of two vaginal Misoprostol regimens, 400 $\mu$ g and 600 $\mu$ g, on pregnancy termination in the 2<sup>nd</sup> trimester

Niroomanesh SH.(M.D.)<sup>1</sup>, Hashemi Fesharaki M.(M.D.)<sup>2</sup>.

1- Professor, Department of Obs & Gyn, Mirza Koochak Khan Hospital, Faculty of Medicine, Tehran University of Medical Sciences, Tehran, Iran.

2- Resident, Department of Obs & Gyn, Mirza Koochak Khan Hospital, Faculty of Medicine, Tehran University of Medical Sciences, Tehran, Iran.

### Abstract

**Introduction:** In a number of complications including intrauterine fetal death, premature rupture of membrane, chorioamnionitis and preeclampsia, gynecologists have to terminate the pregnancy in the 2<sup>nd</sup> trimester to save the mother's life. The methods of pregnancy termination in a uterus larger than 14 weeks in size, include spontaneous contractions, PGE<sub>1</sub> and PGE<sub>2</sub> vaginal suppositories, intravenous oxytocin injection, intra amniotic injection of PGF<sub>2 $\alpha$</sub> , Ethacrynic Acid, and hypertonic saline. Misoprostol, a synthetic PGE1 analogue, which was developed for the prevention of peptic ulcer disease with potent uterotonic and cervical ripening activity, is nowadays applied in the management of obstetric problems like termination of pregnancy. The goal of the present clinical trial is to compare the effectiveness, side effects, hospitalization period and the need for surgical evacuation, when 400  $\mu$ g and 600  $\mu$ g Misoprostol is applied to terminate a second trimester pregnancy.

**Materials and Methods:** One hundred women with medical indications for termination of pregnancy in the second trimester were recruited in Mirza Koochak Khan Hospital in 2002- 2003. They were randomly divided into two groups (each group with 50 cases). One group received 400  $\mu$ g vaginal Misoprostol suppositories (Group A) and the other group 600  $\mu$ g (Group B) every 12 hours in posterior fornix, upto a maximum total of 4 doses. For Comparison of the results in the two groups  $\chi^2$  and t- test were used.

**Results:** In the group A, 84%, as compared to 80% in the group B, had a complete expulsion of the product of conception. Fourteen percent in group A and 20% in group B needed curettage. The success rate of treatment (complete expulsion of pregnancy products) after 48 hours was 98% and 100% in Group A and group B, respectively. The interval between induction and expulsion of fetus was shorter in the group B (9.2 $\pm$ 6.7 hours) than in the group A (12.2 $\pm$ 7.7 hrs, p=0.003) and in patients with intrauterine fetal death (7.4 $\pm$ 2.8 hrs) compared with premature rupture of membrane (11.9 $\pm$  9.6 hrs, p=0.005). The placenta was expelled after the fetus with shorter interval in group A (p=0.02). There was no statistically significant difference between the two groups concerning the duration of hospitalization and maternal complications such as nausea, vomiting, diarrhoea, fever, unrelenting abdominal pain and hemoglobin drop.

**Conclusion:** Vaginal administration of Misoprostol is a quite safe, efficient and cost effective method for termination of pregnancy in the second trimester. In emergency situations such as chorioamnionitis and preeclampsia, 600  $\mu$ g dose might be recommended for rapid expulsion of products of conception, otherwise 400  $\mu$ g dose is a less costly alternative.

**Key Words:** Misoprostol, Second trimester, Premature Rupture of Membrane, and Intrauterine Fetal Death.

**Corresponding Address:** Dr. Niroomanesh, Obs& Gyn Dep., Mirza Koochak Khan Hospital, North Ostad Nejatollahi St., Karimkan-Zand St., Tehran, Iran.

**E mail:** dr.niromanesh@yahoo.com