

Producing Recombinant mTEX101; a Murine Testis Specific Protein

Barzegar Yarmohammadi, Leila (M.Sc.)^{1,2}; Modarresi, Mohammad Hossein (Ph.D.)^{2,3}; Talebi, Saeed (M.D.)^{1,2}; Hadavi, Reza (M.Sc.)¹; Ostad Karampour, Mahyar (M.Sc.)¹; Mahmoudi Ahmad Reza (M.Sc.)¹; Akhondi Mohammad Mehdi (Ph.D.)⁴; Rabbani, Hodjattallah (Ph.D.)^{1,5}; Jeddi-Tehrani, Mahmood (Ph.D.)^{1,5*}

1- Monoclonal Antibody Research Center, Avicenna Research Institute, ACECR, Tehran, Iran.

2- Department of Medical Genetics, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran.

3- Pasteur Institute of Iran, Tehran, Iran.

4- Reproductive Biotechnology Research Center, Avicenna Research Institute, ACECR, Tehran, Iran.

5- Immune and Gene Therapy Laboratory, Karolinska Cancer Center, Karolinska University Hospital, Stockholm, Sweden.

Abstract

Introduction: Production of antibodies against specific proteins of testis germ cells is of great significance for the investigation of processes involved in spermatogenesis, study of infertility problems and determination of the probable role of these proteins as cancer-testis antigens. Murine Testis Specific Recombinant Protein 101 (mTEX101) is a 38kDa, GPI-anchored protein which is expressed in testis germ cells of adult mice but it seems to be absent in other tissues. The structure and function of mTEX101 is not completely understood yet, but it is speculated that it may transduce biochemical signals into the cytoplasm since mTEX101 does not have an intracellular domain but the precise mechanisms are still ambiguous.

Materials and Methods: RNA was extracted from three adult mice testis. The RNA was used in RT-PCR, employing a pair of specific primers for mTEX101 ORF region. TA-cloning technique was performed by the insertion of mTEX101 into a pGEM-T Easy Vector, followed by its subcloning into a His-tagged expression vector, pET-28a (+). The recombinant mTEX101 was then produced by transfection of the expression vector into BL 21 (DE3) E. coli strain.

Results: A recombinant protein, weighing 27kDa, was produced upon IPTG-induction of the bacterial host. The presence of mTEX101 protein was detected through Western blot analysis by anti-mTEX101 peptide antibodies.

Conclusion: We produced mTEX101 recombinant protein that could be used for the production of mono and polyclonal antibodies.

Keywords: Antibody production, Gametogenesis, Germ cell, Glycoprotein, Recombinant protein, Spermatogenesis, TES101 protein mouse, Testis.

To cite this article: Barzegar Yarmohammadi L, Modarresi MH, Talebi S, Hadavi R, Ostad Karampour M, Mahmoudi AR, et al. Producing Recombinant mTEX101; a Murine Testis Specific Protein. *J Reprod Infertil.* 2009;10(3):193-8.

*Corresponding Author:

Dr. Mahmood Jeddi-Tehrani, Monoclonal Antibody Research Center, Avicenna Research Institute, Shahid Beheshti University, Velenjak, P.O. Box: 19615-1177, Tehran, Iran.

E-mail:

mahjed@yahoo.com
mahjed@avicenna.ac.ir

Received: Feb. 3, 2009

Accepted: Jun.15, 2009

Introduction

Spermatogenesis is a complex process of development that occurs in most mammals throughout adulthood period. This process happens in seminiferous tubules, lined with epithelial tissue that contains Sertoli cells. Sertoli cells are

surrounded by a thin septum of peritubular cells that hold keep germ cells (1,2).

The spermatogonia undergo mitotic divisions, meiotic alternations, and morphologically transform to highly developed cells, spermatozoa, with

a unique function and structure (3). Such a complex process needs a unique program to control and regulate the involved genes. This gene regulation is accomplished by cell surface molecules which participate in cell to cell and cell to extracellular matrix interactions (4,5). On the other hand, when sperm is passing through the male and female genital tracts, it gains fertilization ability (Capacitation) and adopts maturity characteristics. Maturity is initiated in the epididymis, where sperm communicates with the cell surface molecules of the pathway (6). Therefore, the investigation of stage and/or cell-specific molecular expression in testis or post-testicular maturation system can improve our understanding of germ cell differentiation and sperm formation (7).

Kurita *et al.* first identified a 38kDa protein (mTEX101) in adult mouse testis which is detectable on the surface of spermatocytes and spermatids in testis that it is absent on somatic cells, including Sertoli or interstitial cells, like Leydig cells (3). MTEX101 shows sexually dimorphic expression during gametogenesis. In the developing ovary, there is a temporary expression of mTEX101 on oogonia, but the molecule disappears in the mature ovary. In embryonic testis, mTEX101 is detected on the prospermatogonia (8). Upon puberty, mTEX101 mRNA is expressed in spermatocytes and in steps 1-9 spermatids during spermatogenesis, but not during spermatogonia. The TEX101 protein remains on the cell surfaces in steps 10-16 spermatids and testicular sperm, but hides from epididymal sperm after passing the caput epididymis (6).

MTEX101 homologues have been identified in rats and humans. Aside from testis, the expression of mTEX101 homologous protein has been reported in a leukemia cell line and lung cancer in rats and humans, respectively (9,10).

The exact structure and function of mTEX101 is not completely understood yet, but it has been speculated that this protein is associated with several transmembrane proteins like Ly6k (11) and Cellubrevin (12) that transduce the extracellular signals to the intracellular molecules. The present study was undertaken to produce recombinant mTEX101, which might pave the

way for the production of specific antibodies for further structural and functional characterization of the molecule.

Materials and Methods

RT PCR: Total RNA was isolated from male gonadal organs (four to 6 week-old mice from Pasteur Institute of Iran). RNA concentration was measured by a BioPhotometer (Eppendorf, Hamburg, Germany) at 260nm. One microgram of the total RNA was reversetranscribed to cDNA by using 200U of Molony Murine Leukemia Virus (RTM-MULV) reverse transcriptase enzyme (Fermentas, Vilnius, Lithuania), and 20pmol of random hexamer primers (Cybergene, Stockholm, Sweden). PCR amplifications for mTEX101 transcript was performed in a volume of 25µl using 10-20ng of testis cDNA, forward and reverse primers (10pmol each), specific for mTEX101 transcript, 10X PCR buffer (2.5µl), dNTP mixture (0.2mM each), 1mM MgCl₂, and 1 unit of Taq DNA polymerase (Roche, Mannheim, Germany). PCR oligonucleotide primers were designed based on the mTEX101 ORF. Optimization of temperature and MgCl₂ were performed to obtain a high quality PCR product. The best result was observed at 60 °C with 1mM MgCl₂. PCR reactions were carried out using a thermal cycler (Eppendorf Mastercycler Gradient) as indicated below: a pre-heating cycle at 95 °C for 3min, 35 cycles of denaturation at 95 °C for 30sec, annealing at 60 °C for 30sec, extension at 72 °C for 60sec, and finally a 7-min cycle at 72 °C. The forward and reverse mTEX101 ORF primers used (Forward: 5'CCG AAT TCA TGG GAG CCT GCC GCA TCC AG 3', Reverse: 5' AGG GAA GTG GGT GAG GGG GGA GCA GAG CGG CCG C 3'), contained EcoRI and NotI restriction enzyme (RE) sites, respectively.

The PCR products were resolved by ethidium bromide-stained 1.5% agarose gel electrophoresis, which revealed a single distinct band at 750bp (Figure 1). The 750bp band was purified using QIA quick Gel Extraction Kit (QIAGEN, Germantown, MD, USA).

Cloning of mTEX101 in TA vector: The pGEM-T Easy Vector (Promega, Madison, WI, USA) was used for the cloning of the purified PCR product.




Figure 1. PCR optimization on mouse testis cDNA by MgCl₂ gradients. 1-4: MgCl₂ concentrations from 1 to 4 mM, respectively, 5: 1Kb DNA ladder, 6: negative control (no DNA).

Ligation reaction was set up by 50ng of pGEM-T Easy Vector (Promega), three units of T4 DNA ligase, 75ng of mTEX101 purified fragment, and 6μl of Rapid 2X Ligation Buffer (Promega). The mixture was incubated at 16 °C for 16hrs. Competent cells of *E. coli* JM109 strain were used for transformation through heat shock method (17). The transformed bacteria were left for 1 hr in LB broth at 37 °C for recovery, and later 100μl of the transformation culture was plated onto an ampicillin (100mg/ml) (Sigma, Louis, MO, USA), IPTG (Sigma) (0.5mM) and X-gal (Sigma), (80μg/ml) containing LB agar plate and was cultured for 16hrs at 37 °C.

White colonies were screened by 25 cycles of colony PCR under the aforementioned conditions. The recombinant plasmids were isolated from confirmed colonies by miniperp kit (QIAGEN) and then digested by EcoRI and NotI restriction enzymes. Plasmid DNA (800ng) was used in 25μl of the total volume, including NotI (15units), (Invitrogen, Carlsbad, CA, USA), EcoRI (15 units), (Invitrogen), and 2.5μl of 10 X Reaction buffer 3 (Invitrogen) and incubated for 1.5 hr at 37 °C. The mTEX101 fragment in pGEM-T Easy was extracted from agarose gel and then subcloned into a pET-28a (+) expression vector (Merck, Darmstadt, Germany) and was digested by the same restriction enzymes above. Ligation reaction was performed by 125ng of digested pET-28a (+), 0.5pmol of mTEX101 purified fragments, 2μl of Rapid 2X ligation buffer (Pro-

mega), and 3 units of T4 DNA ligase (Promega) in 10μl of the total volume and it was incubated overnight at 4 °C. Ligation products were used for transformation of *E. coli* JM109 strain by heat shock method (17). After a recovery time of 1 hour in LB broth, the transformed bacteria were cultured on LB agar and Kanamycin (50μg/ml) (Sigma, USA) containing plates for 16 hrs at 37 °C. The obtained colonies were screened by colony PCR and one of them was confirmed by DNA sequencing. Recombinant plasmid was purified and was then used for transformation of BL21 (DE3), which is a protease deficient strain of *E. coli*.

Recombinant protein production: A single transformed colony was inoculated into 50ml of LB broth, including Kanamycin (50μg/ml) (Sigma, USA) and untransformed BL21 (DE3) bacteria were cultured into 50 ml of LB medium. Incubation was performed under shaking at 37 °C until the optical density (OD) at 600nm reached 0.6. Ten milliliters of each sample was removed as an uninduced control. IPTG (Sigma) was added to the remainder with the final concentration of 1mM and incubation continued for 2-3 more hours. The cells were harvested by centrifugation at 5000g for 5min at 4 °C. Cell lysates were prepared by sonication of pellets in PBS buffer including 1% protease inhibitor (Roche).

Immunoblotting: Protein concentration of the cell lysates were determined by Bradford Protein Assay. Protein solutions were obtained from bacteria by boiling them in a sample buffer that contained Tris-HCl (pH=6.8, 0.5M), SDS (10% W/V), Glycerol (50% V/V), and Bromo Phenol Blue (0.5% W/V), for 5min. Samples (50μl of each) were separated by SDS-PAGE (8%) (Bio-Rad, Hercules, CA, USA) under non-reducing conditions. Untransformed and uninduced bacteria were used as negative controls.

The protein constituents were then electrophoretically blotted onto a polyvinylidene difluoride (PVDF) membrane (Millipore, Billerica, MA, USA), as described by Towbin *et al.* (13). The membrane was blocked in 5% skim milk / 0.1% Tween 20 in PBS (pH 7.4), and then the reactivity of the transferred protein(s) with 1μg/ml




Figure 2. Colony PCR on transformed JM109 clones by pGEM-T Easy vector carrying mTEX101 gene. 1-5: 5 selected white colonies, 6: negative control (blue colony), 7: positive control (PCR product on testis cDNA), 8: DNA ladder VIII (Roche).

of anti mTEX101 pAb (Avicenna Research Institute, Iran) was assessed using sheep anti rabbit Ig HRP (Avicenna Research Institute, Iran). Finally, the membranes were visualized using an ECL system (GE Healthcare, Biotech, Buckinghamshire, UK).

Results

mTEX101 Cloning: After cloning mTEX101 fragments in pGEM-T Easy Vector, several white colonies with probable target fragment inclusion were screened by colony PCR (Figure 2). A right-sized PCR product and a pET-28a(+) vector were cut by NotI and EcoRI restriction enzymes to




Figure 4. Double digestion of pET-28a (+) expression vector with restriction enzymes. 1: digested pET-28a (+) with EcoRI and NotI, 2: uncut pET-28a (+), 3: 1Kb DNA ladder.




Figure 3. Double digestion of pGEM-T easy vector containing mTEX101 fragment with EcoRI and NotI restriction enzymes. 1: digested vector with mTEX101 750bp insert cut out of the vector, 2: DNA ladder VIII.

obtain the required fragments for the next step (Figures 3 and 4).

E. coli JM109 strains were transformed using recombinant pET-28a(+) containing mTEX101 fragment and the undertaken transformation was verified by colony PCR (Fig 5). One of the confirmed colonies (clone 2 in figure 5) was picked for detailed analysis by DNA sequencing. Alignment of DNA sequencing results with mTEX101 ORF by Genbee site (14) confirmed the cloned sequence. The verified fragment was used for the next round of transformation using *E. coli* BL21 (DE3) strain.

Immunoblotting: Western blot analysis using




Figure 5. Colony PCR on transformed JM109 clones by pET-28a (+) vector containing mTEX101 fragment. 1 and 2 represent two selected colonies, 3: DNA ladder VIII, 4: negative control (no DNA), 5: positive control (pGEM-T Easy vector containing mTEX101 fragment).




Figure 6. Western blot analysis of production of the recombinant mTEX101 in BL21 (DE3) bacteria. 1: BL21 (DE3) containing mTEX101 gene after induction, 2: BL21 (DE3) containing mTEX101 gene before induction, 3: Untransfected BL21 (DE3), 4: Protein marker (See Blue-Invitrogen).

anti-mTE X 101 peptide polyclonal antibody revealed the correct size of mTEX101 recombinant protein production (27kDa) in BL-21 (DE3) bacteria (Figure 6).

Discussion

MTEX101 was first reported and characterized by Kurita *et al.* in 2001 (3). The researchers immunized female mice with 8-week old testis lysates and obtained 12 monoclonal antibody producing clones (TES101-112). Immunohistochemistry analyses revealed that only TES101 was able to recognize a novel testicular protein as determined by searching the Expressed Sequence Tag (EST) database (3). All studies on mTEX101 protein are performed solely by TES101 antibody and presently, no commercial antibodies are available to carry out research on this protein. In this study, we produced mTEX101 recombinant protein that is useful for antibody development for further investigations.

MTEX101 mRNA contains 750 bases, which encodes 250 amino acids. After the removal of the putative 25-amino acid signal peptide at N-terminus, the molecular mass of the remaining protein is expected to be about 24kDa. Kurita *et al.* reported a 38kDa band by Western blot analysis under non-reducing conditions and no 24 kDa band was observed (3). This large difference in the molecular mass was speculated to be due to

glycosylation of the peptide and it was proved to be true by Jin *et al.* observations (15).

Glycosylation is known as the prime cause of post-translational modifications (PTM) in proteins (16). The Asn-X-Ser/Thr is the basic sequence for N-linked glycosylation, however, the secondary structure of the protein can affect the final addition. No consensus sequence has been established for O-linked glycosylation yet (17).

Jin *et al.* found four putative sites for N-glycosylation and several possible sites for O-glycosylation in mTEX101 amino acid sequence (15). In fact, mTEX101 is a highly glycosylated protein and it has been clarified that most of the oligosaccharide chains on this peptide are N-linked carbohydrates (12). De-N-glycosylation of mTEX101 created an ~20kDa band which is close to or even smaller than estimated molecular weight of mTEX101 amino acid backbone (15), confirming glycosylation as the cause for the higher mass of native mTEX101.

The cellular role of glycoprotein sugar components can be investigated via several methods, like protein production in a host lacking oligosaccharide addition system, such as prokaryotes (18). We succeeded to produce mTEX101 recombinant protein in a prokaryotic system, which may be useful in clarifying the role of sugar components in protein structures. However, proteins which are produced by this method may vary in their structures and functions (19). Western blot analysis showed that this protein had a molecular mass of approximately 27kDa under non-reducing conditions that is concordant with the various post-translational modification processes in prokaryotic systems as compared to those in eukaryotes. The size of the recombinant protein (27kDa) is in conformity with the backbone protein (24kDa) and a 3kDa peptide derived from the digested vector by NotI and EcoRI restriction enzymes.

Conclusion

In this study, we successfully cloned mTEX101 into a His-tagged expression vector (pET-28a(+)), followed by an efficient production of the relevant recombinant protein. This protein can be used in antibody production to find out the role of

mTEX101 in spermatogenesis, egg fertilization and further investigation about the critical role of glycosylation in the function of this protein.

Acknowledgement

We are deeply grateful to Mrs. R. Ghods, from Avicenna Research Institute, for her encouragements throughout the course of this study. This work was supported by a grant (number:957-33) from Avicenna Research Institute.

References

- Mackay S. Gonadal development in mammals at the cellular and molecular levels. *Int Rev Cytol.* 2000; 200:47-99.
- Starz-Gaiano M, Lehmann R. Moving towards the next generation. *Mech Dev.* 2001;105(1-2):5-18. Review.
- Kurita A, Takizawa T, Takayama T, Totsukawa K, Matsubara S, Shibahara H, et al. Identification, cloning, and initial characterization of a novel mouse testicular germ cell-specific antigen. *Biol Reprod.* 2001;64(3):935-45.
- Fujisawa M. Cell-to-cell cross talk in the testis. *Urol Res.* 2001;29(3):144-51.
- Siu MK, Cheng CY. Dynamic cross-talk between cells and the extracellular matrix in the testis. *Bioessays.* 2004;26(9):978-92.
- Dacheux JL, Gatti JL, Dacheux F. Contribution of epididymal secretory proteins for spermatozoa maturation. *Microsc Res Tech.* 2003;61(1):7-17. Review.
- Takayama T, Mishima T, Mori M, Ishikawa T, Takizawa T, Goto T, et al. TEX101 is shed from the surface of sperm located in the caput epididymidis of the mouse. *Zygote.* 2005;13(4):325-33.
- Takayama T, Mishima T, Mori M, Jin H, Tsukamoto H, Takahashi K, et al. Sexually dimorphic expression of the novel germ cell antigen TEX101 during mouse gonad development. *Biol Reprod.* 2005;72(6):1315-23.
- Hálová I, Dráberová L, Dráber P. A novel lipid raft-associated glycoprotein, TEC-21, activates rat basophilic leukemia cells independently of the type 1 Fc epsilon receptor. *Int Immunol.* 2002;14(2):213-23.
- Tajima K, Obata Y, Tamaki H, Yoshida M, Chen YT, Scanlan MJ, et al. Expression of cancer/testis (CT) antigens in lung cancer. *Lung Cancer.* 2003;42(1):23-33.
- Yoshitake H, Tsukamoto H, Maruyama-Fukushima M, Takamori K, Ogawa H, Araki Y. TEX101, a germ cell-marker glycoprotein, is associated with lymphocyte antigen 6 complex locus k within the mouse testis. *Biochem Biophys Res Commun.* 2008;372(2):277-82.
- Tsukamoto H, Yoshitake H, Mori M, Yanagida M, Takamori K, Ogawa H, et al. Testicular proteins associated with the germ cell-marker, TEX101: involvement of cellubrevin in TEX101-trafficking to the cell surface during spermatogenesis. *Biochem Biophys Res Commun.* 2006;345(1):229-38.
- Towbin H, Staehelin T, Gordon J. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc Natl Acad Sci USA.* 1979;76(9):4350-4.
- GeneBee [internet]. Moscow: A.N. Belozersky Institute of Physico-Chemical Biology; 1965-2009. GeneBee - Molecular Biology Server; 2008 Dec 20 [cited Dec 25]. Available from: http://www.genebee.msu.ru/services/malign_reduced.html
- Jin H, Yoshitake H, Tsukamoto H, Takahashi M, Mori M, Takizawa T, et al. Molecular characterization of a germ-cell-specific antigen, TEX101, from mouse testis. *Zygote.* 2006;14(3): 201-8.
- Campbell CT, Yarema KJ. Large-scale approaches for glycobiology. *Genome Biol.* 2005;6 (11):236. Review.
- Hang HC, Bertozzi CR. The chemistry and biology of mucin-type O-linked glycosylation. *Bioorg Med Chem.* 2005;13(17):5021-34. Review.
- Shriver Z, Raguram S, Sasisekharan R. Glycomics: a pathway to a class of new and improved therapeutics. *Nat Rev Drug Discov.* 2004;3(10): 863-73.
- Walsh G, Jefferis R. Post-translational modifications in the context of therapeutic proteins. *Nat Biotechnol.* 2006;24(10):1241-52. Review.