

Reversible Germ Cell Toxicity of Sulphasalazine and Ampicillin Combination in Male Rats

Himanshu Gupta^{1*}, Kamal Kishore Maheshwari², Nitin Kumar¹

1- Department of Pharmaceutical Technology, Meerut Institute of Engineering and Technology, Uttar Pradesh, India

2- Department of Pharmacy, Mahatma Jyotiba Phule Rohilkhand University, Uttar Pradesh, India

Abstract

Background: The antifertility effect of ampicillin (AMP, 40 mg/kg) and sulphasalazine or salicylazosulfapyridine (SASP, 300,450 and 600 mg/kg) in male rats has been reported earlier. The combination of AMP and SASP is generally used in certain pathological conditions, but the combined effect of these two drugs on the fertility is not clear. So, the aim of this study was to investigate the antifertility effect of ampicillin and sulphasalazine combination in male rats.

Methods: In the present study, forty rats were randomly divided into five groups (n=8). Group I served as the control, while Group II and III received AMP and SASP at the doses of 20 mg/kg and 200 mg/kg respectively. Moreover, group IV and V received the combination of SASP (100 mg/kg) and AMP (10 mg/kg). However, for evaluating the reversible effect of the combination, a washout period of 30 days was given in group V. After 45 days of drug treatment, each rat was sacrificed. The testes, seminal vesicles and epididymis were dissected & weighed. Furthermore, fertility tests, sperm characteristic analysis, histopathological studies, testosterone assay and tissue biochemistry were performed. The data were analyzed using ANOVA and in case ANOVA shows statistical differences, post hoc analysis was performed.

Results: A decrease in parameters related to fertility of males such as sperm count, sperm motility, fertility ratio, serum testosterone level, glycogen and protein content in sexual organs was observed. Although AMP and SASP significantly ($p < 0.001$) reduced the reproductive activity separately, but their combination was found to be impairing the reproductive activity at a considerably lower dose. However, on withdrawing the treatment, all these parameters were restored which was confirmed by the histopathological analysis of the testis.

Conclusion: The combination produces synergistic antifertility effect in male rats and the effect was reversible. The dose and efficacy of results could be extrapolated in future clinical trials.

Keywords: Ampicillin, Fertility, Serum testosterone assay, Sperm count, Sulphasalazine

To cite this article: Gupta H, Maheshwari KK, Kumar N. Reversible Germ Cell Toxicity of Sulphasalazine and Ampicillin Combination in Male Rats. *J Reprod Infertil.* 2013;14(3):126-132.

* Corresponding Author:
Himanshu Gupta,
Department of Pharmacy,
S.R.M.S. College of Engineering and Technology
Bareilly (U.P.), India
E-mail:
himanshu.bpharm@gmail.com

Received: May 11, 2013

Accepted: Jul. 13, 2013

Introduction

Today, the world population is around 7.11 billion and that of India in particular is around 1.23. One of the major problems of the developing countries like India is its geometrical increase in the human population. This population explosion will have negative impact on our economic policies and would simultaneously misbalance our socio-economic infrastructure. Thus, the control of human fertility in the sense of its limitation is the most important and urgent re-

lution explosion will have negative impact on our economic policies and would simultaneously misbalance our socio-economic infrastructure. Thus, the control of human fertility in the sense of its limitation is the most important and urgent re-

quirement (1). So, various scientists have started to solve this problem by developing the contraceptives. Basically contraceptives refer to those chemical agents that either inhibit the sperm motility and sperm production in males or prevent the generation of ovum and produce some structural changes within the endometrium, making it non-receptive to a fertilized ovum in the case of females (2). At recent, several potential approaches for induction of infertility have been investigated over a long period, including chemical, hormonal, and immunological approaches, recently. However, no suitable method has emerged that is effective and free from side-effect (3).

Infertility is defined as the state in which a couple wanting a child cannot conceive even after the 12 months of unprotected sexual intercourse (4). In men, the main causes of infertility are oligospermia, asthenospermia, teratozoospermia and azoospermia, which account for 20–25% of cases (5). There are a number of risk factors such as STD involving *N. gonorrhoeae* and *C. trachomatis*. It leads to changes in semen quality and chronic infection may lead to a block of the vas deferens or seminal vesicles (6). Besides these factors, many drugs may lead to infertility in male. The drugs causing infertility in males are alkylating agents, antibiotics such as ampicillin, dicloxacillin, calcium channel blockers, sulphasalazine, chloroquine and many more (7). The major target sites for fertility regulation in the male reproductive tract are testis, where spermatogenesis and sperm production occur. In epididymis, spermatozoa acquire progressive motility and fertilizing capacity or sperm maturation. This organ represents an ideal extragonadal site for fertility regulation and vas deferens is a passage for the transport of spermatozoa during ejaculation. Any intervention at this site would result in either azoospermia or inability of spermatozoa to initiate fertilization associated events (8).

Cytotoxic drugs cause oligospermia by directly affecting the germinal epithelium. Alkylating agents have a dose-related effect and may cause testicular atrophy, but the function may be regenerated many months after stopping the treatment. Methotrexate given for psoriasis produces oligospermia, though plasma hormone concentrations remain normal, the oligospermia improves within a few months of stopping the drug. Similarly, uses of various sulpha drugs like sulphasalazine were found to be producing varying degrees of infertili-

ty in males. The side effects produced in this way are not unexpected, but more surprising is the influence of SASP. The mechanism of the side effect of using SASP is not clear, but it seems that either SASP or a metabolite is toxic for developing spermatozoa (9). Psychotropic drugs, especially monoamine oxidase inhibitors, are thought to affect the sperm count adversely (7). Similarly, adverse effects of many antibiotics and chloroquine on male reproductive functions have been reported. Nitrofurantoin, macrolides such as erythromycin and tylosin, aminoglycosides such as gentamycin and neomycin (10), tetracyclines, ampicillin and sulfadruugs have all been reported to cause varying degrees of spermatogenic inhibition in men and various animal models (11).

Many researchers have reported the antifertility effect of AMP and SASP at a considerable higher dose (12, 13). Both drugs are used in combination for various pathological conditions like in diarrhoea (14). Therefore, this motivated researchers to use a combination of these drugs for evaluating their synergistic effect after repeated exposure in a mammalian test system.

Methods

Animals: Forty colony bred adult male wistar rats, 5–6 months old and weighing between 175 to 250 gr with proved fertility were used for the study. All the animal experiment protocols were approved by the Institutional Animal Ethics Committee (IAEC) and the experimentation on animals was done in accordance with the CPCSEA (Committee for the Purpose of Control and Supervision of Experimentation on Animals, Approval No.711/02/a/CPCSEA) guidelines. Experiments were performed on male wistar rats procured from the central animal facility of the institute. All the animals were kept under controlled environmental conditions at room temperature ($22\pm 2^\circ\text{C}$) with humidity ($50\pm 10\%$) and a 12 hr light and dark cycle. Standard laboratory animal feed (purchased from commercial supplier) and water were given ad libitum. Animals were acclimatized to experimental conditions prior to the start of dosing for a period of one week.

Chemicals: Sulphasalazine, Ampicillin, Hematoxylin and Eosin, were purchased from Sigma-Aldrich Chemicals, Saint Louis, MO, USA. Ethylenediamine-tetraacetic acid and Hank's balanced salt solution (HBSS) were obtained from HiMedia Laboratories Ltd, Mumbai. Testosterone

estimation kit was purchased from Cayman Chemical, Michigan USA (Testosterone EIA Kit Item Number 582701).

Experimental design: It is reported that SASP at the doses of 300, 450 and 600 mg/kg induced significant decrease in fertility of male rats (13). Raji et al. (12) showed reduction in fertility of male when treated with AMP at the dose of 40 mg/kg. Based on the above studies, for evaluating the synergistic effects of SASP and AMP combination, they were used at the doses of 100 mg/kg and 10 mg/kg respectively. However, for studying their individual effect, 200 mg/kg SASP and 20 mg/kg AMP were used so as to evaluate their anti-fertility effect at a lower dose in comparison to the earlier studies. Apart from it, for studying the reversible effect of the combination, another group received the combination of both drugs for the full treatment period followed by a recovery period of 30 days. The suspension of both drugs was prepared in corn oil and administered through oral route up to a treatment period of 45 days. The volume of administration of drugs to each animal was 10 ml/kg of body weight. All the animals were divided into five groups (n=8). Spermatogenesis is a highly organized process and in rodents, sperm are produced from the progenitor spermatogonia after a series of meiotic and mitotic divisions, which takes approximately 42–56 days (15–17). This explains the basis of taking the sampling after 45 days of exposure. Group I served as the control, while Group II and III received AMP and SASP at the doses of 20 mg/kg and 200 mg/kg respectively. Moreover, group IV and V received the combination of both of them at the doses of 100 mg/kg (SASP) and 10 mg/kg (AMP). However, for evaluating the reversible effect of the combination, a washout period of 30 days was given in group V. Animals were sacrificed by cervical dislocation after the last treatment.

Autopsy and organ weights: At the end of the treatment period, each rat was sacrificed by cervical dislocation. The testes, seminal vesicles and epididymis were dissected, freed from adherent tissue and weighed accurately up to milligram level.

Fertility test: Successful mating (male female ratio 1:2) was carried out with all the animals, five days prior to sacrifice period. The successful mating was confirmed in the following mornings by vaginal plug and spermatozoa in the vaginal

smear. The inseminated females were separated and after gestation period, the number of females delivered, number of litter born and fertility percentage were recorded (8).

Sperm characteristic analysis: After sacrificing the animal, a 1 mm incision was made in the caudal epididymis and drops of sperm fluid were squeezed onto the microscope slide and 2 drops of normal saline were added to mobilize the sperm. Epididymal sperm motility was then assessed by calculating motile spermatozoa per unit area and was expressed in percentage. Epididymal sperm counts were also done by homogenizing the epididymis in HBSS. Counting was then done using the counting chamber in the haemocytometer (18). For sperm head morphology, the sperm suspension in HBSS was stained with 2% eosin solution and kept undisturbed for 1 hr. Smears were prepared using the above solution, air dried and fixed with absolute methanol for 5 min. Two hundred sperm per animal were examined to determine the morphological abnormalities. Sperm head morphology was scored under the category of normal, sperm without hook, amorphous head, banana head and triangular head.

Histopathological studies: After sacrificing the rats, the testes were fixed in 10% formalin, dehydrated in increasing concentrations of ethanol and then embedded in paraffin. Tissue sections (5 µm) were mounted on glass slides which was already coated with Mayer's albumin and dried overnight. The sections were then de-paraffinized with xylene, rehydrated with alcohol and water. The rehydrated sections were stained, mounted with DPX mounting media and examined under the microscope (19).

Testosterone assay: Serum testosterone assay was carried out using the enzyme immunoassay (EIA) method. The within assay variation was 8.1 % and the sensitivity was 0.3 ng/mL. For the estimation of testosterone, blood samples were collected by retro-orbital plexus and serum samples were separated by standard procedures and stored at –20 °C for subsequent analysis.

Tissue biochemistry: The testis and epididymis were dissected out, freed from adherent tissues and weight accurately. Protein (20) and glycogen (21) concentration were estimated in testis and other accessory reproductive organs.

Statistical analysis: Results were shown as Mean ± standard error of mean (S.E.M.) for each group. Statistical analysis was performed using Jandel

Sigma Stat (Version 2.03) statistical software. For multiple comparisons, One-way analysis of variance (ANOVA) was used. In case ANOVA showed significant differences, post hoc analysis was performed with Tukey's test. The level of significance of the data was considered $p < 0.05$.

Results

Effect on weight of body and sex organs: Significant decrease in the final body weight ($p < 0.001$) was observed after 45 days of treatment with the combination of SASP and AMP at the doses of 100 mg/kg and 10 mg/kg as compared with their initial weight (Table 1). The relative weights of different sexual organs were also determined. The weight of testis and seminal vesicle in all treatment groups showed significant difference ($p < 0.001$) after the treatment period of 45 days as compared to control group but in recovery group, no significant changes were observed after 30

days of withdrawing the treatment.

Fertility: The ratio between delivered and inseminated females (09/16, 03/16, 01/16 and 15/16 animals versus 16/16 animals in group I) and the number of pups dropped after treatments are shown in table 2. Although there was a decrease in ratio between delivered and inseminated females in all the groups except for the recovery group but this ratio was found to be least in the group which was treated with the combination of the drugs. However, all the delivered pups were normal and healthy.

Effect on spermatozoa indices: The effects on sperm motility and counts were shown in table 3.

Motility: Significant decrease in sperm motility was found in the group treated with AMP at the dose of 20 mg/kg ($p < 0.05$) as compared with the control count. However, in all the other groups, there was not a significant decrease in sperm motility of the animals.

Table 1. Effects of drug treatments after 45 days on body, testicular, epididymal and seminal vesicle weights in wistar rats

Treatment Group	Body weight (gr)		Testis (gr)	Epididymis (gr)	Seminal vesicle (gr)
	Initial	Final			
Group I (Control)	251±18.19	253±17.12	1.98±0.05	0.43±0.03	0.41±0.02
Group II (AMP 20 mg/kg)	231±16.12	236±16.12	0.91±0.07	0.33±0.05	0.26±0.06*
Group III (SASP 200 mg/kg)	245±17.51	250±12.12	0.79±0.11	0.29±0.04	0.25±0.01*
Group IV (Combination)	255±18.78	196±21.12	0.54±0.08	0.21±0.06	0.19±0.05*
Group V (Recovery)	260±12.19	265±16.22	1.88±0.06	0.33±0.03	0.43±0.09

Values are expressed as Mean±SEM (n=8), * $p \leq 0.01$ when compared with control group

Table 2. Effects of drug treatments on number of females delivered/ number of inseminated females, total number of pups, litter weight and mean percentage fertility (Male: female ratio, 1:2)

Treatment groups	Mean percentage fertility (%)	Total N of pups	Litter weight (gr)
Group I (Control)	100.00	97	9.3±0.91
Group II (AMP 20 mg/kg)	56.25	56	8.1±0.82
Group III (SASP 200 mg/kg)	37.50	51	6.1±1.21
Group IV (Combination)	6.25	04	7.0±1.31
Group V (Recovery)	93.75	94	8.6±1.19

Table 3. Effect of drug treatments after 45 days on sperm motility and sperm count in wistar rats

Treatment Group	Sperm motility (%)	Sperm counts ($\times 10^6/ml$)
Group I (Control)	84.81±2.31	62.00±8.76
Group II (AMP 20 mg/kg)	74.62±1.31*	50.67±9.82**
Group III (SASP 200 mg/kg)	88.98±3.21	52.88±7.61**
Group IV (Combination)	80.22±2.78	29.65±6.54***
Group V (Recovery)	83.21±1.93	61.00±7.67

values are expressed as Mean±SEM (n=8), * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.001$ when compared with control group




Figure 1. Representative photomicrographs of sperm. A: Normal sperm, B: sperm without hook, C: banana head sperm, D: sperm with triangular heads

Epididymal sperm count and sperm head morphology: Decline in sperm count was observed in all the groups. A significant decrease in sperm count was observed ($p < 0.01$) in the groups which were treated with AMP and SASP at the doses of 20 mg/kg and 200 mg/kg respectively as compared to the control group. Furthermore, a significant decrease in sperm count of the animals treated with the combination of AMP and SASP at the doses of 10 mg/kg and 100 mg/kg ($p < 0.001$) was also observed as compared to the control group; however, no significant change in sperm count was seen in recovery group. Apart from this fact, abnormalities in sperm head were seen in all the groups as compared with the control group (Figure 1).

Serum testosterone levels: The mean serum testosterone level was found to be significantly ($p < 0.001$) reduced in all the treatment groups as compared to the control group. However, in the recovery group there was not a significant change in the serum testosterone level (Table 4).

Glycogen and protein levels: The group treated with AMP at the dose of 20 mg/kg showed significant ($p < 0.01$) reduction in glycogen content of both testis and epididymis as compared to the control group. Similarly, group IV which was treated with the combination of both AMP and SASP at the doses of 10 mg/kg and 100 mg/kg showed significant ($p < 0.001$) decrease in glycogen content of both testis and epididymis as compared to that of the control group. Furthermore, the protein levels in both testis and epididymis were significantly reduced in all the groups as compared to that of the control group. There was not a significant change in the recovery group (Table 5).

Histological changes: The histology of both AMP and SASP treated rat testis showed maturation arrest of spermatozoa as compared to control group in which all the stages of spermatogenesis can be seen. But the effect produced by the AMP was patchy. However, in the group which was treated with the combination of both the drugs, the seminiferous tubules showed lack of spermatozoa along with the necrosis of the germ cells. Also, there was disorganization of plasmalemma in the basal portion of few seminiferous tubules. But the histological slide of the recovery group showed all the stages of spermatogenesis (Figure 2).

Table 4. Effects of drug treatments after 45 days on serum testosterone level in wistar rats

Treatment Group	Serum testosterone level (ng/dL)
Group I (Control)	256±1.9
Group II (AMP 20 mg/kg)	90.87±1.2 *
Group III (SASP 200 mg/kg)	69.23±2.1 *
Group IV (Combination)	70.12±1.9 *
Group V (Recovery)	254±2.1

Values are expressed as Mean±SEM (n=8), * $p < 0.001$ when compared with control group

Table 5. Effects of drug treatments after 45 days of treatment on glycogen and protein level in reproductive organs of wistar rats

Treatment Group	Glycogen Level (µg/mg tissue)		Protein level (µg/mg tissue)	
	In testis	In epididymis	In testis	In epididymis
Group I (Control)	5.78±0.15	2.76±0.12	8.92±0.99	4.71±0.1
Group II (AMP 20 mg/kg)	4.01±0.74 *	1.91±0.22 *	6.65±0.65 **	2.43±0.2 **
Group III (SASP 200 mg/kg)	5.34±0.54	1.85±0.56 *	6.43±0.43 **	2.86±0.54 **
Group IV (Combination)	2.87±0.77 **	1.22±0.54 **	6.28±0.33 **	1.31±0.2 **
Group V (Recovery)	5.81±0.82	2.69±0.68	8.88±0.77	4.67±0.3

Values are expressed as Mean±SEM (n=8), * $p < 0.01$, ** $p < 0.001$ when compared with control group




Figure 2. Photomicrographs of stained histological slides of the testis after 45 days of treatment; A: Group I (Control), B: Group II (AMP 20 mg/kg), C: Group III (SASP 200 mg/kg), D: Group IV (Combination), E: Group V (Recovery)

Discussion

The results of this investigation reveal that the administration of SASP, AMP and their combination interfere with the structure and function of major elements of male fertility as reflected by the marked reduction in percentage fertility, serum testosterone level, sperm count and decrease in weight of major sex organs but a significant decrease in motility of sperm was not observed except for the group which was treated with AMP. Similar results were reported by Chumpol Pholpramool et al. (13) for SASP at the doses of 300, 450 and 600 mg/kg and by Y. Raji et al. (12) for AMP at the dose of 40 mg/kg in male rats.

The decreased fructose and protein level may affect glycoproteins secreted by the epididymis and coated on the sperm to stimulate motility (22). The reduced protein content may be another reason for the reduction in the weight of reproductive organs, because the growth rate of organ is proportional to its protein content (23). The depletion in the glycogen content due to treatment hampers the glycolytic metabolism of spermatozoa. A decrease in the glycogen content of the testis reduces the energy source for spermatogenic activity (24). The present study reveals reduction in the level of glycogen and protein of testes and epididymis, and it affects spermatogenesis and sperm maturation.

However, in this investigation, AMP and SASP either separately or in combination were producing decline in the reproductive capacity of male rats but the combination of both SASP and AMP

was found to be impairing the reproductive activity at a considerably lower dose. Apart from it, overall decrease in the fertility, sperm count and serum testosterone level was very higher in comparison to other groups. So, it can be predicted that they have synergistic antifertility effect in male rats. Till now, the exact mode of action of both SASP and AMP as antifertility agents is not well elucidated yet. But according to Marmor et al. (6), SASP produces antifertility action due to inhibition of dietary folic acid absorption, while AMP produces this effect due to inhibition of sperm motility (12).

Conclusion

Thus, our results clearly demonstrate the treatment by AMP and SASP combination in male rats induced synergistic germ cell toxicity in rats as is evident from the decrease in sperm count, serum testosterone level and decline in percentage fertility but the effect was reversible as is evident from the recovery group. Furthermore, this was confirmed by histopathological analysis and sperm head morphology but the cause of this infertility at genetic level is not evaluated in the research paper. As they are both used in combination for treatment of various pathological conditions, it warrants further attention for complete elucidation of the reasons behind this toxic effect so as to reduce the potential risk in the patients, who are particularly in the reproductive ages.

Acknowledgement

The financial assistance received from department of pharmaceutical technology, MIET Meerut to carry out this particular study is duly acknowledged.

Conflict of Interest

Authors declare no conflict of interest.

References

1. Kamal R, Gupta RS, Lohiya NK. Plants for male fertility regulation. *Phytother Res.* 2003;17(6):579-90.
2. Ziarati P. Determination of contaminants in some Iranian popular herbal medicines. *J Env Anal Toxicol.* 2012;2:120.
3. Ma WK, Ramaswamy SB. Histological changes during ovarian maturation in the tarnished plant bug, *Lygus lineolaris* (Hemiptera: Miridae). *J Insect Morphol Embryol.* 1974;16:304-32.

4. Olayemi FO. A review on some causes of male infertility. *Afr J Biotech*. 2010;9:2834-42.
5. Egozcue S, Blanco J, Vendrell JM, García F, Veiga A, Aran B, et al. Human male infertility: chromosome anomalies, meiotic disorders, abnormal spermatozoa and recurrent abortion. *Hum Reprod Update*. 2000;6(1):93-105.
6. Marmor D. The effects of sulphasalazine on male infertility. *Reprod Tox*. 1995;9:219-23.
7. Drife JO. Drugs and sperm. *Br Med J (Clin Res Ed)*. 1982;284(6319):844-5.
8. Shivabasavaiah KRH, Pavana T, Ramyashree M, Ramya MC, Manjunath R. Antifertility effects of madhuca indica leaves in male swiss albino rats. *J Pharm Res*. 2011;4:323-6.
9. Pholpramool C, Srikhao A. Suppression of fertility by sulfasalazine and its metabolite in the male rats. *J Androl*. 1985;6:50.
10. Timmermans L. Influence of antibiotics on spermatogenesis. *J Urol*. 1974;112(3):348-9.
11. Kushniruk YI. Influence of certain antibacterial preparations on the nucleic acid content in cells of the spermatogenic epithelium. *Tsitol Genet*. 1976;10:342-7.
12. Raji Y, Awobajo FO, Alabi TK, Gbadegsin MA, Bolarinwa AF. In vivo and in vitro reproductive toxicity assesment of ampicillin and cloxacillin in mammalian models. *Int J Pharmacol*. 2006;2:9-14.
13. Pholpramool C, Srikhao A. Antifertility effect of sulfasalazine in the male rat. *Contraception*. 1983;28(3):273-9.
14. Tripathi KD. *Essential of Medical Pharmacology*. 5th ed. India: Jaypee Brothers Medical Publishers; 2004. 754p.
15. Peirce EJ, Breed WG. A comparative study of sperm production in two species of Australian arid zone rodents (*Pseudomys australis*, *Notomys alexis*) with marked differences in testis size. *Reproduction*. 2001;121(2):239-47.
16. Olsen AK, Lindeman B, Wiger R, Duale N, Brunborg G. How do male germ cells handle DNA damage? *Toxicol Appl Pharmacol*. 2005;207(2 Suppl):521-31.
17. Aragón MA, Ayala ME, Fortoul TI, Bizarro P, Altamirano-Lozano M. Vanadium induced ultrastructural changes and apoptosis in male germ cells. *Reprod Toxicol*. 2005;20(1):127-34.
18. Padmanabhan S, Tripathi DN, Vikram A, Ramarao P, Jena GB. Cytotoxic and genotoxic effects of methotrexate in germ cells of male Swiss mice. *Mutat Res*. 2008;655(1-2):59-67.
19. Adeeko AO, Dada OA. Chloroquine reduces fertilizing capacity of epididyma sperm in rats. *Afr J Med Med Sci*. 1998;27(1-2):63-4.
20. Lowry OH, Rosenbrough NJ, Far AL, Randall RJ. Protein measurement with the Folin phenol reagent. *J Biol Chem*. 1951;193(1):265-75.
21. Montgomery R. Determination glycogen. *Arch Biochem Biophys*. 1957;67(2):378-86.
22. Gupta G, Rajalakshmi M, Prasad MR, Moudgal NR. Alteration of epididymal function and its relation to maturation of spermatozoa. *Andrologia*. 1974;6(1):35-44.
23. Sarkar M, Gangopadhyay P, Basak B, Chakrabarty K, Banerji J, Adhikary P, et al. The reversible anti-fertility effect of Piper betle Linn. on Swiss albino male mice. *Contraception*. 2000;62(5):271-4.
24. Rommerts FF, Cooke BA, van der Molen HJ. The role of cyclic AMP in the regulation of steroid biosynthesis in testis tissue. *J Steroid Biochem*. 1974;5(3):279-85.