

The Effect of Curcumin on Intracellular pH (pHi), Membrane Hyperpolarization and Sperm Motility

Rajesh K. Naz

- Reproductive Immunology and Molecular Biology Laboratories, Department of Obstetrics and Gynecology, West Virginia University, School of Medicine, Morgantown, West Virginia, USA

Abstract

Background: Curcumin has shown to affect sperm motility and function *in vitro* and fertility *in vivo*. The molecular mechanism(s) by which curcumin affects sperm motility has not been delineated. Since modulation of intracellular pH (pHi) and plasma membrane polarization is involved in sperm motility, the present study was conducted to investigate the effect of curcumin on these sperm (human and murine) parameters.

Methods: The effect of curcumin on sperm forward motility was examined by counting percentages of forward moving sperm. The effect of curcumin on intracellular pH (pHi) was measured by the fluorescent pH indicator 2,7-bicarboxyethyl-5,6-carboxyfluorescein-acetoxymethyl ester (BCECF-AM). The effect of curcumin on plasma membrane polarization was examined using the fluorescence sensitive dye bis (1,3-dibarbituric acid)-trimethine oxanol [DiBAC₄(3)].

Results: Curcumin caused a concentration-dependent ($p < 0.05$) decrease in forward motility of both human and mouse sperm. It also caused a concentration-dependent decrease in intracellular pH (pHi) in both human and mouse sperm. Curcumin induced significant ($p < 0.05$) hyperpolarization of the plasma membrane in both human and mouse sperm.

Conclusion: These findings indicate that curcumin inhibits sperm forward motility by intracellular acidification and hyperpolarization of sperm plasma membrane. This is the first study to our knowledge which examined the effect of curcumin on sperm pHi and membrane polarization that affect sperm forward motility. These exciting findings will have application in deciphering the signal transduction pathway involved in sperm motility and function and in development of a novel non-steroidal contraceptive for infertility.

Keywords: Contraception, Curcumin, Polarization, Signal transduction, Sperm forward motility, Sperm.

To cite this article: Naz RK. The Effect of Curcumin on Intracellular pH (pHi), Membrane Hyperpolarization and Sperm Motility. *J Reprod Infertil.* 2014;15(2):62-70.

* Corresponding Author:
Rajesh K. Naz, Robert C.
Byrd Health Sciences
Center North, West Virginia
University, School of
Medicine, Room 2085,
1 Medical Center Drive,
Morgantown, WV 26506,
USA
E-mail:
Rnaz@hsc.wvu.edu

Received: Dec. 10, 2013

Accepted: Apr. 12, 2014

Introduction

Curcumin, 1,7-bis(4-hydroxy-3-methoxyphenyl)-1,6-heptadiene-3,5-dione, commonly known as diferuloylmethane, is the yellow pigment component of the curry or turmeric (*Curcuma longa*) (1). Turmeric extracts have been extensively used for the treatment of several diseases in Ayurvedic medicine in India for several cen-

turies. Curcumin was first extracted from turmeric in its impure form in 1815, but it was not until 1910 when it was crystallized and its structure was elucidated (2). It has antimicrobial, antioxidant, immunomodulatory, anti-inflammatory, anti-Alzheimer and anticancer activity (3-8). It has shown no toxicity *in vitro* in numerous cell culture

systems, and *in vivo* in animal models and over 13 Phase I human clinical trials. It is generally recognized as safe by the United States Food and Drug Administration (FDA) (2, 4, 5).

Curcumin has been shown to affect several targets in somatic cells for its biological activity (3-8). It inhibits NF- κ B activity, COX-2, and 5-LOX expression and modulates release of several cytokines (3, 4). It also binds to a number of other proteins including thioredoxin reductase, protein kinases and several receptors (3, 4). However, most of these proteins/factors may not be expressed/present in terminally non-transcriptional sperm. Also, the sperm has a unique characteristic, the motility, that is not present in other cells. Thus, curcumin may have different molecules/mechanism(s) for its action that are unique to sperm.

Recently, our laboratory reported, for the first time ever, that the curcumin affects sperm function (motility/capacitation and acrosome reaction/fertilization) *in vitro* and fertility *in vivo*. Intravaginal administration of curcumin caused a significant, but reversible reduction in fertility (9). The molecular mechanism(s) by which curcumin inhibits/blocks sperm motility has not been delineated. Since modulation of intracellular pH (pHi) and plasma membrane polarization has been shown to be involved in sperm motility and capacitation/acrosome reaction of several mammalian species (10-21), the present study was conducted to investigate the effect of curcumin on sperm intracellular pH and plasma membrane polarization. It was examined using both human and mouse sperm. It was hypothesized that the curcumin-mediated effect on sperm motility is caused by modulation of pHi and/or membrane polarization. The long-term objective of the study was to understand the molecular mechanism(s) by which curcumin affects sperm motility and function and to develop a novel non-steroidal contraceptive with spermicidal properties.

Methods

Collection of sperm: Human spermatozoa were collected from healthy, fertile men. Semen was liquefied and analyzed for volume, sperm concentration and percent and progressive motility. Only those semen samples that had sperm concentration of $>50 \times 10^6$ sperm/ml, percent motility of $>60\%$, progressive motility of $>+3$ (on a scale of 0 to +5), and contamination of immature germ cells and immune cells of $<1\%$ were used to collect a pure swim-up sperm population (22). The study was

approved by the West Virginia University-Institutional Review Board (IRB) for Human Studies.

Mouse sperm were collected from cauda epididymis and vas deferens of mature BALB/c or CD-1 males. Motile sperm were isolated by the swim-up procedure and washed by centrifugation (500 g, 10 min) with Ham's F-10 medium supplemented with human serum albumin (5 mg/ml) or Modified Sperm Washing MediumTM (Irvine Scientific, Santa Ana, CA, USA). Motile sperm were analyzed for sperm concentration and motility (9). The study was approved by the West Virginia University-Animal Care and Use Committee (ACUC) for animal studies.

Measurement of sperm forward motility: The highly purified form of curcumin (cat. #C27727) was obtained from Sigma-Aldrich (St. Louis, MO, USA). Curcumin was dissolved in dimethylsulfoxide (DMSO) (25-50 mM stock) and then diluted in medium to the desired concentrations. Sperm treated with medium or equivalent volume of DMSO served as the controls. The effect of curcumin on sperm forward motility was examined by incubating 10-100 μ l of sperm suspension ($100-250 \times 10^6$ motile sperm/ml) with various concentrations of curcumin (50-400 μ M, final concentration) up to 1 hr. The percentage of forward moving sperm was recorded every 5-20 min, before and after incubation. Sperm treated with an equivalent volume of medium or DMSO served as the controls. Each treatment in each experiment was tested in duplicate. Each experiment, including various treatment groups, was performed 3-5 times using sperm from 3-5 men or mice on different days to take care of the inter-individual variation among various sperm specimens from different men and mice.

Measurement of sperm intracellular pH: The intracellular pH (pHi) of human and mouse sperm was measured by fluorescent pH-indicator 2,7-bis-carboxyethyl-5,6-carboxyfluorescein-acetoxymethyl-ester (BCECF-AM) (Molecular Probes, Eugene, OR, USA) following the manufacturer's protocol as described by Hamamah et al. (23). BCECF is a neutral lipophilic form of bis-carboxyfluorescein which diffuses freely through the plasma membrane. In the cell, it is hydrolyzed by esterases, releasing the BCECF which is retained within the cytoplasm. The fluorescence intensity of BCECF is dependent upon the pH. The motile sperm, isolated by the swim-up procedure, were centrifuged and the pellet was washed and resuspended in 1 ml of phosphate-buffered saline (PBS, pH=7.4).

Sperm ($8-15 \times 10^6/ml$) were then loaded with $2 \mu M$ BCECF (final concentration) and incubated ($37^\circ C$, 35 min) in dark. Following incubation, sperm were centrifuged, washed (x2) and resuspended in PBS. For the intracellular pH calibration curve, sperm were loaded with BCECF at various extracellular pHs (pHe; 6.8, 7.0, 7.2, 7.4, 7.6 and 7.8) and then treated with 0.1% Triton-X100. Subsequently, the fluorescence intensity was measured as described below, and the calibration curve was constructed by plotting fluorescence intensity versus extracellular pH (24). Using calibration curve, the fluorescence intensity value for each treatment was converted to its respective intracellular pH.

The fluorescence intensity of BCECF is dependent upon the pH with a maximum response at an excitation of $\lambda=535 \text{ nm}$ (F1), while at $\lambda=490 \text{ nm}$ (F2), the intensity is independent of pH. The pHi was determined graphically using the ratio F1/F2 from a calibration curve obtained after permeabilization of spermatozoa with 0.1% Triton by measuring the maximum fluorescence intensity after adding NaOH and the minimum after adding HCl. The F1/F2 ratio represents a pseudo-linear function of the pH (23).

To examine the effect of curcumin, BCECF-loaded sperm were incubated ($37^\circ C$, $5-10 \text{ min}$) with various concentrations of curcumin ($50-400 \mu M$) and then washed. The treated/control sperm were transferred into wells ($200 \mu L/well$) of a black 96-well microplate and the fluorescence intensity was measured (BioTek Synergy2 Multiplatform automated plate reader) using excitation and emission wavelengths of 490 and 535 nm , respectively. Each treatment in each experiment was tested in duplicate. Each experiment, including various treatment groups, was performed 3-5 times using sperm from 3-5 men or mice on different days to take care of the inter-individual variation among various sperm specimens from different men and mice.

All of the appropriate controls were carefully included as described by the manufacturer. Moreover, it was pertinent to examine the effect of curcumin per se on fluorescence intensity. Similar experiments carried out with various concentrations ($50-400 \mu M$) of curcumin without sperm did not affect the fluorescence intensity, indicating curcumin per se, without sperm, does not affect fluorescence intensity.

Plasma membrane polarization of human and mouse sperm: The changes in sperm plasma membrane polarization were examined using the fluo-

rescence sensitive dye bis (1,3-dibarbituric acid)-trimethine oxanol [DiBAC₄(3)] (Molecular Probes, Eugene, OR, USA) following the manufacturer's protocol as described by Rossato et al. (25). The motile sperm in the swim-up fraction were centrifuged, washed (x2) with PBS and incubated ($37^\circ C$, 1 hr) with $2 \mu M$ DiBAC₄(3). DiBAC₄(3) can enter depolarized cells and then bind to intracellular proteins or membrane and exhibits enhanced fluorescence. Increased depolarization results in additional influx of the anionic dye and an increase in fluorescence. Conversely, hyperpolarization is indicated by a decrease in fluorescence. This dye is excluded from mitochondria because of their overall negative charge. Sperm ($8-15 \times 10^6/ml$) were then centrifuged and washed twice with PBS. The DiBAC₄(3)-incubated sperm were aliquoted in different tubes at equal volumes and incubated ($5-10 \text{ min}$) with various concentrations of curcumin ($100 \mu M$, $200 \mu M$, $300 \mu M$, and $400 \mu M$). Sperm treated with PBS or equivalent volume of DMSO served as the controls. After final washing with PBS, the sperm were transferred into wells ($200 \mu L/well$) of a black 96-well microplate and the fluorescence intensity was measured (BioTek Synergy2 Multiplatform automated plate reader) using excitation and emission wavelengths of 485 and 530 nm , respectively. Each treatment in each experiment was tested in duplicate. Each experiment, including various treatment groups, was performed 3-5 times using sperm from 3-5 men or mice on different days to take care of the inter-individual variation among various sperm specimens from different men and mice.

All the appropriate controls were carefully included as described by the manufacturer. It was pertinent to examine the effect of curcumin per se on fluorescence intensity. Similar experiments carried out with various concentrations ($50-400 \mu M$) of curcumin without sperm did not affect the fluorescence intensity, indicating curcumin per se, without sperm, does not affect fluorescence intensity.

Statistical analysis: The significance of difference among various groups was analyzed using analysis of variance (ANOVA) and Tukey Kramer Multiple Comparison Test. A $p \leq 0.05$ was considered statistically significant.

Results

Effect of curcumin on human and mouse sperm forward motility: Curcumin caused a concentration-dependent decrease in human sperm forward mo-

tility (Table 1). At 50 μM concentration, there was no apparent ($p>0.05$) effect on sperm forward motility in 5-10 *min* (with a slight decrease over time) and in 1 *hr* it decreased by up to $\sim 25\%$ ($p<0.001$). At 100 μM concentration, there was a significant ($p<0.001$) decrease in sperm forward motility in 5-10 *min* which decreased by up to $\sim 80\%$ in 1 *hr*. At concentrations $\geq 200 \mu\text{M}$, there was a complete block of sperm forward motility within 5-10 *min*. DMSO vehicle did not affect ($p>0.05$) sperm forward motility as compared to medium control.

Curcumin caused a similar effect on mouse sperm forward motility (Table 2). At 50 μM concentration, there was no apparent ($p>0.05$) effect on sperm forward motility in 5-10 *min* (with a slight decrease over time) and in 1 *hr*, it decreased by up to $\sim 40\%$ ($p<0.001$). At 100 μM concentration, there was a significant ($p<0.001$) decrease in sperm forward motility in 5-10 *min*, which decreased by up to $\sim 70\%$ in 1 *hr*. At concentrations $\geq 200 \mu\text{M}$, there was a complete block of forward motility within 5-10 *min*. DMSO vehicle did not affect ($p>0.05$) sperm forward motility as compared to the medium control.

Effect of curcumin on human and mouse sperm intracellular pH (pHi): There was a linear relationship between the extracellular pH and the fluorescence intensity both in human ($R^2=0.9513$) and mouse ($R^2=0.9835$) sperm (Figure 1, panel A and B, respectively). With an increase in extracellular pH, there was a corresponding increase in fluorescence intensity. The $\text{pH}=7.4$ was selected in our subsequent experiments to examine the effect of curcumin on intracellular sperm pH (pHi).

Table 1. Effect of curcumin on human sperm forward motility (%)*

Treatment	0 min	5-10 min	20 min	40 min	60 min
Curcumin					
50 μM	95 \pm 5	91 \pm 6	85 \pm 5	80 \pm 7	72 \pm 8**
100 μM	95 \pm 5	40 \pm 4**	26 \pm 7**	20 \pm 8**	18 \pm 7**
200 μM	95 \pm 5	0 \pm 0**	0 \pm 0**	0 \pm 0**	0 \pm 0**
300 μM	95 \pm 5	0 \pm 0**	0 \pm 0**	0 \pm 0**	0 \pm 0**
400 μM	95 \pm 5	0 \pm 0**	0 \pm 0**	0 \pm 0**	0 \pm 0**
DMSO vehicle					
	95 \pm 5	95 \pm 5	95 \pm 5	95 \pm 5	93 \pm 6
Medium control					
	95 \pm 5	95 \pm 5	95 \pm 5	95 \pm 5	93 \pm 6

* Mean \pm SD; ** Versus control, significantly different ($p<0.001$); all others non-significant

Table 2. Effect of curcumin on murine sperm forward motility (%)*

Treatment	0 min	5-10 min	20 min	40 min	60 min
Curcumin					
50 μM	92 \pm 8	85 \pm 7	80 \pm 6	71 \pm 4**	61 \pm 7**
100 μM	92 \pm 8	42 \pm 9**	35 \pm 4**	30 \pm 6**	28 \pm 6**
200 μM	92 \pm 8	0 \pm 0**	0 \pm 0**	0 \pm 0**	0 \pm 0**
300 μM	92 \pm 8	0 \pm 0**	0 \pm 0**	0 \pm 0**	0 \pm 0**
400 μM	92 \pm 8	0 \pm 0**	0 \pm 0**	0 \pm 0**	0 \pm 0**
DMSO vehicle					
	92 \pm 8	92 \pm 8	92 \pm 8	90 \pm 7	85 \pm 8
Medium control					
	92 \pm 8	92 \pm 8	92 \pm 8	90 \pm 7	88 \pm 7

* Mean \pm SD; ** Versus control, significantly different ($p<0.001$); all others non-significant

In both human and mouse sperm, curcumin caused a concentration-dependent decrease in intracellular pH (Figure 2), in human (panel A) and mouse (panel B) sperm. In human sperm, control sperm had pHi of 7.3 ± 0.003 which was not significantly different ($p>0.05$) from DMSO-treated sperm, which had a pHi of 7.3 ± 0.105 (Figure 3, panel A). Treatment with curcumin significantly ($p<0.001$) decreased intracellular pH in a concentration-dependent manner as compared to control/DMSO-treated sperm. At 50 μM curcumin concentration, pHi was 7.21 ± 0.010 , at 100 μM pHi was 7.04 ± 0.008 , at 200 μM pHi was 6.95 ± 0.02 , at 300 μM pHi was 6.88 ± 0.0067 and at 400 μM pHi was 6.81 ± 0.014 . Comparing the change in pHi within the curcumin-treated groups, the difference was significant ($p<0.05$) only among 50 μM , 100 μM and 400 μM groups.

Similar results were obtained in mouse sperm (Figure 2, panel B). In mouse sperm, control sperm had pHi of 7.15 ± 0.005 which was not significantly different ($p>0.05$) from DMSO-treated sperm which had a pHi of 7.15 ± 0.00 (Figure 2, panel B). Treatment with curcumin significantly ($p<0.001$) decreased intracellular pH in a concentration-dependent manner as compared to control/DMSO-treated sperm. At 50 μM curcumin concentration, pHi was 7.07 ± 0.017 , at 100 μM pHi was 6.97 ± 0.017 , at 200 μM pHi was 6.91 ± 0.012 , at 300 μM pHi was 6.88 ± 0.02 and at 400 μM pHi was 6.83 ± 0.017 . Comparing the changes in pHi within the curcumin-treated groups, the difference was significant ($p<0.05$) only among 50 μM , 100 μM and 400 μM groups.




Figure 1. Calibration curve of human (panel A) and mouse (panel B) sperm fluorescence intensity at various extracellular pH. BCECF-loaded sperm were treated with 0.1% Triton-X 100 at various extracellular pH (pHe) and the fluorescence intensity was measured. Fluorescence intensity values were then plotted against pHe values to obtain the calibration curve.




Figure 2. Human (panel A) and mouse (panel B) sperm intracellular pH (pHi) after curcumin treatment. BCECF-loaded sperm were treated with increasing concentrations of curcumin and the fluorescence intensities were measured and converted to intracellular pH using the calibration curve. Intracellular pH (pHi) values (Mean±SE) from 3-5 independent experiments using sperm from 3-5 different men and mice are shown. There was no effect of any concentration of curcumin tested per se on fluorescence intensity without sperm.

Effect of curcumin on human and mouse sperm plasma membrane polarization: The effect of various concentrations of curcumin on plasma membrane polarization was evaluated using both human and mouse sperm. In human sperm, treatment with DMSO did not significantly ($p>0.05$) affect fluorescence intensity as compared to control sperm (control: 2861 ± 43.016 ; DMSO: 3252 ± 215.860) (Figure 3, panel A). Treatment with curcumin caused significant ($p<0.001$) hyperpolarization of sperm plasma membrane as compared to control/DMSO-treated sperm (Figure 3, panel A). At $50\ \mu\text{M}$ curcumin, fluorescence intensity were 601 ± 51.052 , at $100\ \mu\text{M}$, 597 ± 20.664 , at $200\ \mu\text{M}$, 522 ± 72.421 , at $300\ \mu\text{M}$, 458 ± 47.576 and at 400

μM 414 ± 29.464 . Comparing the changes in fluorescence intensities within the curcumin-treated groups, the differences were not-significant.

Similar results were obtained in mouse sperm (Figure 3, panel B). In mouse sperm, treatment with DMSO did not significantly affect fluorescence intensity as compared to control (control: 5592.085 ± 2.186 ; DMSO: 5130.905 ± 130.770) (Figure 3, panel B). Treatment with curcumin caused significant ($p<0.001$) hyperpolarization of sperm plasma membrane as compared to control/DMSO-treated sperm (Figure 3, panel B). At $50\ \mu\text{M}$ curcumin, fluorescence intensity was 997.502 ± 22.229 , at $100\ \mu\text{M}$ was 877.356 ± 41.328 , at $200\ \mu\text{M}$ was 624.331 ± 42.116 , at $300\ \mu\text{M}$ was 578.929 ± 43.245




Figure 3. Effect of curcumin on plasma membrane potential in human (panel A) and mouse (panel B) sperm. Fluorescence intensity values (Mean \pm SE) from 3-5 independent experiments using sperm from 3-5 different men and mice are shown. Fluorescence intensity values lower than the control indicate hyperpolarization. There was no effect of any concentration of curcumin tested per se on fluorescence intensity without sperm.

and at 400 μ M was 522.431 \pm 57.900. Comparing the changes in fluorescence intensity within the curcumin-treated groups, the difference was significant ($p < 0.05$) only between 50 μ M and 400 μ M groups.

Discussion

The data indicated that curcumin affects sperm forward motility starting at 100 μ M concentration, with a complete block at ≥ 200 μ M concentration within 5-10 *min* in both human and murine sperm. These findings correlate well with the earlier published data (9). In general, forward motility corresponded well with overall motility and progressive motility, as a decrease in forward motility corresponded with decrease in overall and progressive motility. Also, there was a loss of sperm viability when there was a decrease in forward motility. The totally immotile sperm were completely non-viable as tested by eosin-nigrosin staining (data not shown).

The present study was conducted to examine the mechanism by which curcumin affects sperm forward motility. The findings indicated that curcumin acidified sperm intracellularly and hyperpolarized cell membrane of both human and mouse sperm. There was a concentration-dependent decrease in pHi from 7.3 to 6.81 at highest concentration (400 μ M) of curcumin tested. These findings correlate well with the effect of curcumin on human and murine sperm forward motility. Starting at 100 μ M concentration, there was a significant effect within 5-10 *min*, with a total block at ≥ 200 μ M concentration. All the human and murine sperm samples tested demonstrated a similar phenotype. DMSO vehicle did not affect the pHi.

The pHi plays an important role in modulating mammalian sperm motility (12, 17, 20). These cumulative findings indicate that the acidic pHi inhibits sperm motility, which is in agreement with our findings. The pHi of 6.6-6.8 has been reported in one study for uncapacitated murine sperm (13). In this study, the pHi of control murine sperm was slightly higher, ~ 7.15 . This may be due to different methodologies/media and/or the strain of mice used and/or various degrees of capacitation of sperm preparations. The pHi of 7.25 was observed for control human sperm in this study. This is in agreement with a previous published study (24). The sodium-proton membrane exchange mechanism can modulate pHi and affect motility (26). The pHi changes are also involved in sperm motility as the sperm cell passes through the epididymis (27, 28). Also, the pH of the epididymal cauda fluid is acidic in almost all species (29) and acts directly on the pHi to decrease the sperm motility (28). Our extensive data search in PubMed and Google Scholar did not yield any publication which has examined the effect of curcumin on intracellular pH in sperm or any other cell types.

Curcumin caused hyperpolarization of sperm plasma membrane in both human and mouse sperm. The degree of decrease in intracellular fluorescence intensity is indicative of the degree of hyperpolarization. Hyperpolarization is a change which makes cell's membrane potential more negative. It is the opposite of depolarization. The change in sperm membrane potential has been shown to be involved in sperm motility and function (18, 19, 21, 24, 25, 28, 30-33). For mouse sperm, intracellular alkalinization produces a hy-

perpolarization of sperm membrane potential which may be important for both hyperactivated motility and acrosome reaction. In this study, curcumin induced hyperpolarization and intracellular acidification which correlate with a reduction/block of sperm motility. There was more drastic effect on hyperpolarization than on change in pHi. At 50 μM concentration, which did not significantly decrease the pHi, it showed a significant effect on membrane polarization in both human and mouse sperm. It is possible that curcumin may affect pHi and membrane potential by interacting with different molecules and signal transduction cascades. Besides sperm, hyperpolarization has also been connected to intracellular pHi changes in several other cell systems (34).

Although there is no publication which has examined the effect on sperm, curcumin has been reported to affect membrane potential in other cell systems. Curcumin affects plasma membrane permeability of rat thymocytes (35). Curcumin inhibits sarco/endoplasmic reticulum $\text{Ca}^{++}\text{ATPase}$ (Ser CA) pump that has a major role in maintaining lower levels of intracellular $[\text{Ca}^{++}]$ by importing extracellular protons in rabbit skeleton muscle (36). A study by Cao et al. (37) examined the effect of curcumin on changes in mitochondrial membrane potential in human hepatomaG2 (HepG2) cells. Their study showed that HepG2 cells incubated with curcumin induced oxidative damage to mitochondrial DNA through mitochondrial membrane hyperpolarization.

Several molecules/mechanisms are involved in regulation of sperm intracellular pH and membrane polarization. Sodium-hydrogen exchange has been suggested as a mediator of pH regulation in various mammalian sperm (27, 38, 39). At least two sodium-hydrogen exchangers (NHE1 and NHE5) have been shown to be expressed in spermatozoa. The catalytic subunit of protein kinase A (PKA) is involved in activation of motility evoked by HCO_3^- anion (40). Capacitation-associated hyperpolarization involves a decrease in intracellular $[\text{Na}^+]$ regulated by PKA through activation of the cystic fibrosis transmembrane conductance regulator (CFTR) (41). Alkalinization activates the intracellular pH-sensitive I(KSper), inducing the membrane potential to approach negative potentials where Ca^{2+} entry via I(CatSper) is maximized (33, 42). The exact molecular mechanism(s) and signal transduction pathway involved in modulation of sperm intracellular pH and membrane potential by curcumin need further investigation.

There are several mechanisms that can affect sperm motility/function leading to infertility (43, 44). This preliminary data indicates that curcumin may inhibit tyrosine phosphorylation of a subset of sperm surface proteins and Ca^{2+} channels (unpublished data). Tyrosine phosphorylation has been shown to be involved in sperm motility, capacitation/acrosome reaction and function (16).

Conclusion

The findings indicate that curcumin causes intracellular acidification and membrane hyperpolarization which may be involved in inhibiting sperm forward motility. This is the first study to our knowledge that examined the effect of curcumin on sperm pHi and membrane potential. These exciting findings will have application in deciphering the molecular mechanism(s) involved in curcumin action and in delineating the signal transduction pathway relevant to sperm motility and function. Also, this data may have clinical application in development of a novel non-steroid contraceptive with spermicidal properties.

Acknowledgement

We sincerely thank Bidur Paudel, Morgan Lough, Clara Beth Novotny, and Vanessa Cuppett for performing several of these experiments and their help in data analysis and excellent typing assistance. We also thank Don Zeh and Melanie Clemmer, Ph.D., at the Andrology Laboratory at the Center for Reproductive Medicine at West Virginia University for help in consenting men for providing semen specimens used in this study. This research did not receive any specific grant from any funding agency in the public, commercial or not-for-profit sector.

Conflict of Interest

The author reports no conflict of interest.

References

1. Tayyem RF, Heath DD, Al-Delaimy WK, Rock CL. Curcumin content of turmeric and curry powders. *Nutr Cancer*. 2006;55(2):126-31.
2. Singh S. From exotic spice to modern drug? *Cell*. 2007;130(5):765-8.
3. Sharma RA, Gescher AJ, Steward WP. Curcumin: the story so far. *Eur J Cancer*. 2005;41(13):1955-68.
4. Aggarwal BB, Sundaram C, Malani N, Ichikawa H. Curcumin: The Indian solid gold. *Adv Exp Med Biol*. 2007;595:1-75.

5. Corson TW, Crews CM. Molecular understanding and modern application of traditional medicines: Triumphs and trials. *Cell*. 2007;130(5):769-74.
6. Gupta SC, Patchva S, Koh W, Aggarwal BB. Discovery of curcumin, a component of the golden spice, and its miraculous biological activities. *Clin Exp Pharmacol Physiol*. 2013;39:283-99.
7. Yallapu MM, Ebeling MC, Jaggi M, Chauhan SC. Plasma proteins interaction with curcumin nanoparticles: implications in cancer therapeutics. *Curr Drug Metab*. 2013;14(4):504-15.
8. Yallapu MM, Jaggi M, Chauhan SC. Curcumin nanomedicine: a road to cancer therapeutics. *Curr Pharm Des*. 2013;19(11):1994-2010.
9. Naz RK. Can curcumin provide an ideal contraceptive? *Mol Reprod Dev*. 2011;78(2):116-23.
10. Garbers DL, Watkins HD, Hansbrough JR, Smith A, Misono KS. The amino acid sequence and chemical synthesis of speract and speract analogues. *J Biol Chem*. 1981;257(6):2734-7.
11. Babcock DF, Rufo GA Jr, Lardy HA. Potassium-dependent increases in cytosolic pH stimulate metabolism and motility of mammalian sperm. *Proc Natl Acad Sci USA*. 1983;80(5):1327-31.
12. Babcock DF, Pfeiffer DR. Independent elevation of cytosolic [Ca²⁺] and pH of mammalian sperm by voltage-dependent and pH-sensitive mechanisms. *J Biol Chem*. 1987;262(31):15041-7.
13. Zeng Y, Oberdorf JA, Florman HM. pH regulation in mouse sperm: identification of Na⁽⁺⁾-, Cl⁽⁻⁾-, and HCO₃⁽⁻⁾-dependent and arylaminobenzoate-dependent regulatory mechanisms and characterization of their roles in sperm capacitation. *Dev Biol*. 1996;173(2):510-20.
14. Hamamah S, Gatti JL. Role of the ionic environment and internal pH on sperm activity. *Hum Reprod*. 1998 Dec;13 Suppl 4:20-30.
15. Darszon A, Iabarca P, Nishigaki T, Espinosa F. Ion channels in sperm physiology. *Physiol Rev*. 1999;79(2):481-510.
16. Naz RK, Rajesh PB. Role of tyrosine phosphorylation in sperm capacitation/acrosome reaction. *Reprod Biol Endocrinol*. 2004;2:75.
17. Gagnon C, de Lamirande E, editors. Controls of sperm motility. Cambridge: Cambridge University Press; 2006. 108 p. (Jonge CD, Barratt C, editors. *The Sperm Cell*).
18. Zeng XH, Yang C, Kim ST, Lingle CJ, Xia XM. Deletion of the Slo3 gene abolishes alkalization-activated K⁺ current in mouse spermatozoa. *Proc Natl Acad Sci USA*. 2011;108(14):5879-84.
19. De La Vega-Beltran JL, Sanchez-Cardenas C, Krapf D, Hernandez-Gonzalez EO, Wertheimer E, Trevino CL, et al. Mouse sperm membrane potential hyperpolarization is necessary and sufficient to prepare sperm for the acrosome reaction. *J Biol Chem*. 2012;287(53):44384-93.
20. Lishko PV, Kirichok Y, Ren D, Navarro B, Chung JJ, Clapham DE. The control of male fertility by spermatozoan ion channels. *Ann Rev Physiol*. 2012;74:453-75.
21. Chavez JC, de la Vega-Beltran JL, Escoffier J, Visconti PE, Trevino CL, Darszon A, et al. Ion permeabilities in mouse sperm reveal an external trigger for Slo3-dependent hyperpolarization. *PLoS One*. 2013;8(4):e60578.
22. Samuel AS, Naz RK. Isolation of human single chain variable fragment antibodies against specific sperm antigens for immunocontraceptive development. *Hum Reprod*. 2008;23(6):1324-37.
23. Hamamah S, Magnoux E, Royere D, Barthelemy C, Dacheux JL, Gatti JL. Internal pH of human spermatozoa: effect of ions, human follicular fluid and progesterone. *Mol Hum Reprod*. 1996;2(4):219-24.
24. Jain R, Jain A, Kumar R, Verma V, Mikhuri JP, Sharma VL, et al. Functional attenuation of human sperm by novel, non-surfactant spermicides: precise targeting of membrane physiology without affecting structure. *Hum Reprod*. 2010;25(5):1165-76.
25. Rossato M, Di Virgilio F, Rizzuto R, Galeazzi C, Foresta C. Intracellular calcium store depletion and acrosome reaction in human spermatozoa: role of calcium and plasma membrane potential. *Mol Hum Reprod*. 2001;7(2):119-28.
26. Wong PYD, Lee WM, Tsang AYW. The effects of extracellular sodium on acid release and motility initiation in rat caudal epididymal spermatozoa in vitro. *Exp Cell Res*. 1981;131(1):97-104.
27. Carr DW, Acott TS. Intracellular pH regulates bovine sperm motility and protein phosphorylation. *Biol Reprod*. 1989;41(5):907-20.
28. Gatti JL, Billard R, Christen R. Ionic regulation of the plasma membrane potential of rainbow trout (*Salmo gairdneri*) sperm: role in the initiation of motility. *J Cell Physiol*. 1990;143(3):546-54.
29. Setchell BP, Maddocks S, Brooks DE, editors. Anatomy, vasculature, innervation, and fluids of the male reproductive tract. New York: Raven Press; 1994. 1063 p. (Knobil E, Neill JD, editors. *The Physiology of Reproduction*, vol. 1).
30. Brewis IA, Morton IE, Mohammad SN, Browes CE, Moore HD. Measurement of intracellular cal-

- cium concentration and plasma membrane potential in human spermatozoa using flow cytometry. *J Androl.* 2000;21(2):238-49.
31. Patrat C, Serres C, Jouannet P. Induction of a sodium influx by progesterone in human spermatozoa. *Biol Reprod.* 2000;62(5):1380-6.
 32. Patrat C, Serres C, Jouannet P. Progesterone induces hyperpolarization after a transient depolarization phase in human spermatozoa. *Biol Reprod.* 2002;66(6):1775-80.
 33. Navarro B, Kirichok Y, Clapham DE. KSper, a pH sensitive K⁺ current that controls sperm membrane potential. *Proc Natl Acad Sci USA.* 2007;104(18):7688-92.
 34. Gresik M, Kolarova N, Farkas V. Hyperpolarization and intracellular acidification in *Trichoderma viride* as a response to illumination. *J Gen Microbiol.* 1991;137(11):2605-9.
 35. Jaruga E, Salvioli S, Dobrucki J, Chrul S, Bandorowicz-Pikula J, Sikora E, et al. Apoptosis-like, reversible changes in plasma membrane asymmetry and permeability, and transient modifications in mitochondrial membrane potential induced by curcumin in rat thymocytes. *FEBS Lett.* 1998;433(3):287-93.
 36. Bilmen JG, Khan S, Javed MH, Michelangeli F. Inhibition of the SERCA Ca²⁺ pumps by curcumin. Curcumin putatively stabilizes the interaction between the nucleotide-binding and phosphorylation domains in the absence of ATP. *Eur J Biochem.* 2001;268(23):6318-27.
 37. Cao J, Liu Y, Jia J, Zhou HM, Kong Y, Yang G, et al. Curcumin induces apoptosis through mitochondrial hyperpolarization and mtDNA damage in human hepatoma G2 cells. *Free Radic Biol Med.* 2007;43(6):968-75.
 38. Schackmann RW, Chock PB. Alteration of intracellular [Ca²⁺] in sea urchin sperm by the egg peptide speract. Evidence that increased intracellular Ca²⁺ is coupled to Na⁺ entry and increased intracellular pH. *J Biol Chem.* 1986;261(19):8719-28.
 39. Garcia MA, Meizel S. Regulation of intracellular pH in capacitated human spermatozoa by a Na⁺/H⁺ exchanger. *Mol Reprod Dev.* 1999;52(2):189-95.
 40. Morgan DJ, Weisenhaus M, Shum S, Su T, Zheng R, Zhang C, et al. Tissue specific PKA inhibition using a chemical genetic approach and its application to studies on sperm capacitation. *Proc Natl Acad Sci USA.* 2008;105(52):20740-5.
 41. Escoffier J, Krapf D, Navarrete F, Darszon A, Visconti PE. Flow cytometry analysis reveals a decrease in intracellular sodium during sperm capacitation. *J Cell Sci.* 2012;125(Pt 2):473-85.
 42. McPartlin LA, Visconti, PE, Bedford-Guaus SJ. Guanine nucleotide exchange factors (RAPGEF3/RAPGEF4) induce sperm membrane depolarization and acrosomal exocytosis in capacitated stallion sperm. *Biol Reprod.* 2011;85(1):179-88.
 43. Ohl DA, Naz RK. Infertility due to antisperm antibodies. *Urology.* 1995;46(4):591-602.
 44. Sikka SC. Role of oxidative stress and antioxidants in andrology and assisted reproductive technology. *J Androl.* 2004;25(1):5-18.