

Review of *Trichomonas vaginalis* in Iran, Based on Epidemiological Situation

Mohsen Arbabi *, Mahdi Delavari, Zohreh Fakhrieh-Kashan, Hossein Hooshyar

- Department of Medical Parasitology, School of Medicine, Kashan University of Medical Sciences, Kashan, Iran

Abstract

Trichomoniasis, which is caused by *Trichomonas vaginalis*, is the most common non-viral sexually transmitted infection (STI) in the world including Iran. There were roughly 250 million new cases all over the world in a year. *T. vaginalis* as an important disease has been associated with HIV (in terms of exposure to sexually transmitted infection, STI) which increases the number of high-risk members, and thus it is an important public health problem. Additionally, this pathogen has been associated with serious health consequences. For instance, it may cause a woman to deliver a low-birth-weight or premature infant, and increase chances of cervical cancer. Because little information is available about the prevalence of *T. vaginalis* infection in Iranian population, this review was carried out to determine the prevalence of *T. vaginalis* among Iranian population. For this systematic review, data about epidemiology of *T. vaginalis* in different parts of Iran with different populations were systematically collected from 1992 to 2017 through the international databases such as PubMed, Scirus, ISI Web of Science, Scopus, EMBASE, Science Direct and Google Scholar and Islamic World Science Citation Center (ISC). National database searching included Iran Medex, Iran Doc, Magiran and Scientific Information Database (SID). A total of 39 clinical and laboratory investigations about the prevalence of Trichomoniasis from different regions of Iran were analyzed. The overall prevalence rate of *T. vaginalis* infection in Iranian population was estimated to be minimally 0.4% and maximally 42%. The present review showed that *T. vaginalis* infection rate is relatively high among the Iranian population. The control strategies, including personal hygienic education, simultaneous couple treatment, the sensitivity of diagnostic methods, appropriate preventive tool (condom) in sexual contacts could lead to the disruption of transmission.

* Corresponding Author:
Mohsen Arbabi,
Department of Medical
Parasitology and Mycology,
School of Medicine, Kashan
University of Medical
Sciences, Kashan, Iran
E-mail:
arbabi4.mohsen@yahoo.
com

Received: Oct. 27, 2017

Accepted: Feb. 24, 2018

Keywords: Epidemiology, General population, Iran, Prevalence, *Trichomonas vaginalis*.

To cite this article: Arbabi M, Delavari M, Fakhrieh-Kashan Z, Hooshyar H. Review of *Trichomonas vaginalis* in Iran, Based on Epidemiological Situation. *J Reprod Infertil.* 2018;19(2):82-88.

Introduction

Trichomonas vaginalis is the causative agent of trichomoniasis, a non-ulcerative sexually transmitted disease. *T. vaginalis* is the protozoan parasite infecting the urogenital tract of both females and males (1, 2). It is reported to be 250 million new cases all over the world every year (3). Estimates of prevalence are differences between populations, but the range from 5-74% in women and 5-29% in men is observed (1). Women by the age of 16-53 are at greater risk of infection (4-6). Recently, different studies have

shown that *T. vaginalis* has been associated with HIV, which increases the number of high-risk members (7, 8). Additionally, this pathogen has been associated with serious health consequences; including low-birth-weight in pregnant woman or premature infant and increased chances of cervical cancer. Women who are infected can be asymptomatic or have different symptoms, consisting a yellowish-green frothy discharge purities, dysuria, and the strawberry cervix which is recognized by punctuates hemorrhagic lesions. In general, infec-

tion is asymptomatic in men, although it can be associated with urethral discharge and dysuria (1-4). For diagnosis of trichomoniasis, different methods have been used, such as, wet mount, culture, Papanicolaou smear, Pap smear, polymerase chain reaction (PCR) and serological tests. Wet mount tests are quick and straightforward. Specialized medium cultures are used for diagnosis, but 2-5 days are required. Also, in some cases, parasites are diagnosed in the Papanicolaou smear. Moreover, the lack of sensitivity and specificity of serological examines is the major limitation for the detection of *T. vaginalis* by indirect serological testing. Recently, new approaches to diagnosis of parasite infections are provided by molecular biological methods. PCR allows the amplification of DNA fragments and diminishes the probability of misdiagnosis. When few trophozoites are recognized in a man's reproductive organs, it is believable that PCR is useful for the diagnosis of trichomoniasis (6-9). 5'-nitroimidazole family include metronidazole and tinidazole which are the most common drugs for treatment of trichomoniasis (10, 11). Because little information is available about the prevalence of *T. vaginalis* infection in Iranian population, this systematic review was carried out to determine the prevalence of *T. vaginalis* infection among Iranian population from 1992 to 2017 through the electronic databases.

Literature review of prevalence of trichomoniasis: For this systematic review, electronic searches in international and national databases and journals were conducted using key words of *Trichomonas vaginalis*, general population, prevalence, epidemiology, and Iran.

These articles had used at least one method (Direct smear, culture, PCR, Pap smear) for epidemiological study in different parts of Iran. Searches were performed through the international databases such as PubMed, Scirus, ISI Web of Science, Scopus, EMBASE, Science Direct, Google Scholar and Islamic World Science Citation Center (ISC). National database searching included Iran Medex, Iran Doc, Magiran and Scientific Information Database (SID). Articles that were published between 1992 and 2017 were reviewed. Among the numerous information sources, relevant studies on *T. vaginalis* infection were identified. Articles related to women with childbearing age who referred to health centers due to pregnancy care and symptoms of vaginitis as well as gynecologic problems and men in different parts of Iran were included. A total of 39 papers were in-

vestigated and 3 articles related to women prisoners were excluded from the study. Due to high-risk relationships and inappropriate health conditions, the prevalence of vaginal Trichomoniasis in female prisoners is high and is different from normal rate in the society. Therefore, articles published in the prison population were excluded. For this pathogenic parasitic protozoa, the studies were contradictory and generally of poor quality. At present, different methods have been used for recognizing Trichomoniasis, such as culture, Pap smear, polymerase chain reaction (PCR), and direct smear. Thus, analysis of data was based on the diagnosis of *T. vaginalis* parasite in women who referred to health centers. Results were obtained from different geographical areas classified and analyzed by descriptive statistics.

Women referred to health centers in Sirjan (1992), Kashan (1993), Isfahan (1995) and Tabriz (1998) were infected with *T. vaginalis* (Direct smear: 2.2% and culture: 2.8%), (Culture: 20.1%), (Direct smear: 1.49% and culture: 1.92%), (Direct smear: 22.6%), respectively (12-15). Pap smear specimens from women referring to health centers in Yasouj (1999) were 1.9% infected with *T. vaginalis* (16). In 1010 Pap smear specimens (1999) of 9.9% patients referring to Kashan, were infected with *T. vaginalis* (17). Pregnant women referring to health centers in Kashan (2000) and women referring to health centers in Zahedan (2001) and Hamedan (2001) were infected with *T. vaginalis* (Culture: 0.44%), (Direct smear: 4.5% and culture: 5.3%), (Direct smear: negative), respectively (18-20). Result of the direct smear method and culture in pregnant women referring to health centers in Tehran (2002) showed 2.9% of patients were infected with *T. vaginalis* (21). Women referring to health centers in Gorgan (2003), Hamedan (2004) and Orumieh (2004) were infected with *T. vaginalis* (Direct smear and culture: 9%), (Direct smear: 2% and culture: 3%), (Direct smear: 2.4% and culture: 2.6%), respectively (22-24). 63 women with symptoms of vaginitis in Yasouj were surveyed using direct smear and culture methods (2004) and the result showed 19.04% and 42.9% of patients were infected with *T. vaginalis*, respectively (25). In the health center in Robat Karim (2005), patients were tested by direct smear and culture; results showed 1.4% of them were infected with *T. vaginalis* (26). Sharbatdaran et al. used three methods of direct smear, culture and Pap smears for diagnosis of infection with *T. vaginalis* in Babol (2005) that 18.67%,

18.67% and 25.3% of women were infected respectively (27). 0.9% infection with *T. vaginalis* was observed in 33690 Pap smear specimens in Kermanshah (2005) (28). A survey of women referring to health centers in Tabriz (2006), Yazd (2006), Tehran (2007) showed the percentage of infection with *T. vaginalis* using direct smear and culture was direct smear: 3.46 and culture: 4.56, direct smear: 1.2 and culture: 2.6%, direct smear and culture: 4, direct smear: 22.6, respectively (29-31). In a study that carried out among 300 women referring to health centers of Shahrod (2008) using Pap smear and direct smear methods, only one case of *T. vaginalis* infection was reported (32). A survey of Pap smear samples showed 3.2% of women had a Trichomoniasis infection in the Sari health center (33). Study on women referring to Tehran health center (2009) was achieved using direct smear and culture and obtained result showed the percentage of infection was 2.6 and 3.2, respectively (34). The infection rate of *T. vaginalis* in 160 women suspected of Trichomoniasis in Lorestan was 11.8% and 18.75% using direct smear and culture methods, respectively (35). From 1353 Pap smears specimens in Ahvaz (2010), 1.4% of patients were infected with *T. vaginalis* (36). In two studies in 2010 and 2011, percentage of *T. vaginalis* infection in women with childbearing age and pregnant women in Zanjan was 6.4 and 3.3, respectively (37, 38). PCR-SSCP method was used to test 950 samples from Hamadan and Tehran and obtained results showed fifty samples were positive (39). 3,500 women referring to health centers in Tehran and Kashan were tested using direct smear, culture and PCR methods and 4% of them were infected (40). Patients in Kermanshah, Kashan and rural area of Shahrekord were infected with *T. vaginalis* (Direct smear: 1.5% and culture: 2.1%), (Direct smear and culture: 2%) and (Direct smear: 4%), respectively (41-43). In 2014, women referring to Tehran health centers and Qom were infected with *T. vaginalis* (Culture with urine sample of 5% and vaginal sample of 2.4%, PCR with urine sample of 8.2% and vaginal sample of 8.7%) and (Direct smear: 2.67%, PCR:11.3%), respectively (44, 45) (Table 1).

Estimates of prevalence of *T. vaginalis* between populations are different worldwide, but the range from 5-74% in women and 5-29% in men is observed (1). The incidence of Trichomoniasis has increased remarkably especially in developing countries and in populations with high-risk behav-

iors such as poor sexual activity hygiene and multiple sexual partners. Poverty, socioeconomic status, illiteracy, high risk sexual behaviors, and HIV positive are risk factors for infection of *T. vaginalis* (46-49). In some studies, infection with *T. vaginalis* was more in illiterates than literates (12, 13, 18, 27, 39, 42, 44). Low rate of infection with Trichomoniasis was observed among people who used condoms as a contraceptive method (12, 27, 39, 44), whereas in some studies, Trichomoniasis was not related to prevention methods (13, 14, 42). The reports of Trichomoniasis in Iran are different just the same as other parts of the world. The difference in the prevalence of infection may depend on the selection of population groups, methods and the site of specimen collection. Symptomatic Trichomoniasis is less common in men than women. Biological differences between the two sexes are the cause that women have a higher incidence of infection compared to men (50, 42). Sex hormone is a major factor in different prevalence of Trichomoniasis between both sexes (42). Infection in women can be either asymptomatic or symptomatic, while, it is asymptomatic in men (1-4). Diagnosis of Trichomoniasis on the basis of clinical examinations indicates 88% false negative and 29% false positive results (40). On the basis of some studies, clinical symptoms such as burning and itching were associated with Trichomoniasis (12, 14, 18, 22, 24, 27, 39, 44), while these results were in contrast with other studies (13, 29, 42). Direct smear is the most common method for diagnosis of this infection. After taking samples, they must be tested quickly because the parasite stops moving in a short period and will cause false negative results. The standard method in diagnosis of Trichomoniasis is culture. As this method is sensitive, appropriate conditions including ingredients of culture media, culture temperature, incubation time and the rapid transmission of the parasite after sampling to culture medium are required. Before reporting negative results, negative specimens should be kept up to 7 days for more evaluation. But this method is not used as a routine diagnosis since it wastes much time (44). Typical symptomatic Trichomoniasis in men is cleared spontaneously within 10 days. On the other hand, infection in women can persist for years. Therefore, recognition of carriers is very important for accelerating treatment and decreasing the spread of the disease in control strategies (48, 49). One of the most sensitive diagnostic techniques is polymerase chain reaction (PCR).

Table 1. Prevalence of *T. vaginalis* in different parts of Iran from 1992 to 2017

City	Method	Study population	Year	Prevalence
Sirjan	Direct smear and culture	500 women referred to health centers	1992	Direct smear: 2.2%, Culture: 2.8%
Kashan	Culture	900 women referred to health centers	1993	Culture:2.1%
Esfahan	Direct smear and culture	470 women referred to health centers	1995	Direct smear:1.49%, Culture:1.92%
Tabriz	Direct smear	469 women referred to health centers	1998	Direct smear: 22.6%
Yasouj	Pap smear	1942 Pap smear cytology samples section	1999	Pap smear:1.9%
Kashan	Pap smear	1010 Pap smear cytology samples section	1999	Pap smear: 9.9%
Kashan	Culture	450 pregnant women referred to health centers	2000	Culture:0.44%
Zahedan	Direct smear and culture	597 women referred to health centers	2001	Direct smear:4.5%, culture:5.3%
Hamedan	Direct smear	31 women admitted to the psychiatric ward	2001	Direct smear: negative
Tehran	Direct smear and culture	<i>Pregnant women referred to hospital</i>	2002	Direct smear and culture: 2.9%
Gorgan	Direct smear and culture	102 women referred to health centers	2003	Direct smear and culture: 9%
Hamedan	Direct smear and culture	400 women referred to health centers	2004	Direct smear:2%, culture:3%
Orumieh	Direct smear and culture	420 women referred to health centers	2004	Direct smear:2.4%, culture:2.6%
Yasouj	Direct smear and culture and clinical signs	63 women with symptoms of vaginitis	2004	Clinical signs:19.04%, direct smear and culture:42.9%
Robot Karamu	Direct smear and culture	500 women referred to health centers	2005	Direct smear and culture:1.4%
Babol	Direct smear and culture and Pap smear	150 women with clinical signs	2005	Direct smear:18.67%, culture:18.67%, Pap smear: 25.3%
Kermanshah	Pap smear	33690 Pap smear specimens in clinic	2005	Pap smear: 0.9%
Tabriz	Direct smear and culture	2630 women referred to health centers	2006	Direct smear:3.46%, Culture: 4.56%
Yazd	Direct smear and culture	384 women referred to health centers	2006	Direct smear:1.2%, Culture: 2.6%
Tehran	Direct smear and culture	150 women referred to health centers	2007	Direct smear and culture:4%
Shahrod	Pap smear and direct smear	300 women referred to health centers	2008	Pap smear and direct smear: Only 1
Sari	Pap smear	1832 Pap smear in women referred to health centers	2008	Pap smear: 3.2%
Tehran	Direct smear and culture	500 women referred to health centers	2009	Direct smear: 2.6% , Culture: 3.2%
Lorestan	Direct smear and culture	160 women suspected of Trichomoniasis	2010	Direct smear:11.8%, Culture:18.75%
Ahvaz	Pap smear	1353 Pap smear in women referred to health centers	2010	Pap smear:1.4%
Zanjan	Direct smear	328 women referred to health centers	2010	Direct smear: 6.4%
Zanjan	Direct smear and culture	1000 pregnant women	2011	Direct smear and culture:3.3%
Hamadan and Tehran	PCR-SSCP	950 women referred to health centers	2011	PCR-SSCP: Fifty <i>T. vaginalis</i> samples
Kashan and Tehran	Direct smear , culture and PCR	3500 women referred to health centers	2011	Direct smear , culture and PCR:4%
Kermanshah	Direct smear and culture	600 women referred to health centers	2011-2012	Direct smear1.5%, culture:2.1%
Kermanshah	Direct smear and culture	600 women referred to health centers	2012	Direct smear:1.5%, culture:2.1%
Kermanshah	Pap smear and direct smear	1100 women referred to health centers	2006-2012	Pap smear and direct smear: 0.63%
Kashan	Direct smear and culture	970 women and 235 men referred to health centers	2013	Direct smear and culture :2%
Shahrekord	Direct smear	92 rural women	2014	Direct smear:4%
Tehran	Culture and PCR	140 women referred to health centers	2014	Culture: urine sample: 5%, vaginal sample: 2.4%, PCR: urine sample: 8.2% , vaginal sample: 8.7%
Qom	Direct smear and PCR	300 women referred to health centers	2014	Direct smear: 2.67%, PCR:11.3%
Ardabil	Direct smear and culture	904 women referred to health centers	2014	Direct smear: 3.38%, culture: 4.48%
Hamadan	Direct smear and culture	1200 women referred to health centers	2015	Direct smear: 0.3%, culture: 0.6%
Hamadan	Direct smear and culture	862 women referred to health centers	2015	Direct smear and culture:1.9%

For better chances of accurate diagnosis, at least two techniques are needed, such as wet mount microscopy and culture. Although PCR is found to be highly specific and sensitive, it is costly to be used in routine diagnostic laboratories (42, 49, 51). Based on reviewing research conducted in Iran, studies have used at least one detection method (Pap smear, direct smear, culture and PCR) (41-55). Among the articles, only 12 articles had used a detection method while the other remaining articles used at least 2 or 3 detection methods (12-45, 52-55). Research done among various groups of women indicates that the prevalence is different between women admitted to the psychiatric ward and women with symptoms of vaginitis from zero to 42.9%, respectively (21, 26). Results of this systematic review shows the considerable rate of Trichomoniasis in Iran is related to lack of appropriate control programs in different parts of the country. Control of *T. vaginalis* could have considerable public health benefits in controlling both HIV and sexual diseases in women. This approach needs to be conducted in large communities, with particular attention to high-risk groups. Screening or empiric treatment is not only needed in high-risk groups, but also is needed in low-risk groups like pregnant women.

Conclusion

The present review showed that *T. vaginalis* infection rate is relatively high among the Iranian population. Range of control strategies, including personal hygienic education, simultaneous couple treatment, the sensitivity of diagnostic methods, appropriate preventive tool (condom) in sexual contacts could lead to the disruption of transmission. Using at least two techniques such as culture or PCR in addition to direct smear is recommended for better diagnosis of infection and understanding the actual prevalence of *T. vaginalis*.

Conflict of Interest

We declare that there is no conflict of interest regarding the publication of this paper.

References

- Vos T, Flaxman AD, Naghavi M, Lozano R, Michaud C, Ezzati M, et al. Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012;380(9859):2163-96.
- Mao M, Liu HL. Genetic diversity of *Trichomonas vaginalis* clinical isolates from Henan province in central China. *Pathog Glob Health*. 2015;109(5):242-6.
- Kashan ZF, Arbabi M, Delavari M, Hooshyar H, Taghizadeh M, Joneydy Z. Effect of *Verbascum thapsus* ethanol extract on induction of apoptosis in *Trichomonas vaginalis* in vitro. *Infect Disord Drug Targets*. 2015;15(2):125-30.
- Menezes CB, Mello Mdos S, Tasca T. Comparison of permanent staining methods for the laboratory diagnosis of trichomoniasis. *Rev Inst Med Trop Sao Paulo*. 2016;58:5.
- Mahmoud A, Sherif NA, Abdella R, El-Genedy AR, El Kateb AY, Askalani AN. Prevalence of *Trichomonas vaginalis* infection among Egyptian women using culture and Latex agglutination: cross-sectional study. *BMC Womens Health*. 2015;15:7.
- Khatoun R, Jahan N, Ahmad S, Khan HM, Rabbani T. Comparison of four diagnostic techniques for detection of *Trichomonas vaginalis* infection in females attending tertiary care hospital of North India. *Indian J Pathol Microbiol*. 2015;58(1):36-9.
- Manshoori A, Mirzaei S, Valadkhani Z, Kazemi Arababadi M, Rezaeian M, Zainodini N, et al. A diagnostic and symptomatological study on Trichomoniasis in symptomatic pregnant women in Rafsanjan, south central Iran in 2012-13. *Iran J Parasitol*. 2015;10(3):490-7.
- Secor WE, Meites E, Starr MC, Workowski KA. Neglected parasitic infections in the United States: trichomoniasis. *Am J Trop Med Hyg*. 2014;90(5):800-4.
- Ryu JS, Min DU. *Trichomonas vaginalis* and trichomoniasis in the Republic of Korea. *Korean J Parasitol*. 2006;44(2):101-16.
- Forbes GL, Drayton R, Forbes GD. A case of metronidazole-resistant *Trichomonas vaginalis* in pregnancy. *Int J STD AIDS*. 2015;27(10):906-8.
- Hawkins I, Carne C, Sonnex C, Carmichael A. Successful treatment of refractory *Trichomonas vaginalis* infection using intravenous metronidazole. *Int J STD AIDS*. 2015;26(9):676-8.
- Shaifi I, Khatami M, Tahmores-Kermani E. [Prevalence of *Trichomonas vaginalis* in women referred to vali-asr polyclinic and the health center number 3 in Sirjan city]. *J Kerman Univ Med Sci*. 1994;1(3):125-32. Persian.
- Rasti S, Arbabi M, Khakbazan S, Khamechian T, Hooshyar H, Yadegarifard G. [Epidemiology of Trichomoniasis in women referring to health and therapeutic centers of Kashan in 1372 and 1373]. *Feyz*. 2000;3(4):104-10. Persian.

14. Baghaei M, Memarzadeh Z. [Prevalence of Trichomoniasis in women: Isfahan 1995]. *J Res Med Sci.* 2001;6(2):108-12. Persian.
15. Shahbazi A, Falah E, Safaian A. [Infection rate of Trichomonas vaginalis in females referring to Tabriz Health Care centers, 1998-99]. *Res Med.* 2001; 25(4):231-4. Persian.
16. Parhizkar S, Moshfeh A. [Prevalence of cervicovaginal infections among the Pap smears of women (Yasuj, 1999 -2000)]. *Armaghan Danesh.* 2003;7 (28):37-44. Persian.
17. Rasti S, Khamechian T. [Frequency cytological alterations Trichomoniasis in symptomatic females referring to a gynecology clinic in Kasha]. *Feyz.* 2004;8(1):73-7. Persian.
18. Rasti S, Taghriri A, Behrashi M. [Trichomoniasis in parturient referring to Shabikhani hospital in Kasha, 2001-2002]. *Feyz.* 2003;7(2):21-5. Persian.
19. Zangiabadi M, Qureshi M, Khoushideh M, Roudbari M, Bahrami Sh. Survey of sensitivity of wet smear and dorset medium in comparison with diamond medium for diagnosis of Trichomonas vaginalis. *Zahedan J Res Med Sci.* 2002;4(3):9-15.
20. Tavakol P, Zahirnia AH, Sardarian K, Nazari M, Taherkhani HA, Siavashi MR, et al. [A study of fungal and parasitic infections of skin, digestive and reproductive tract in patients with chronic psychiatric disorders at Sina Hospital in Hamadan (2002-3)]. *J Ilam Univ Med Sci.* 2006;14(3):45-51. Persian.
21. Akbarian A, Akhlaghi L, Ourmazdi H, Foroohesh H, Falahati M, Farokhnejad R. [An investigation on coincidence of Trichomoniasis and bacterial vaginosis and their effects on pregnant women referred to Shahid akbarabadi maternity hospital in Tehran during 2002-2003]. *Razi J Med Sci.* 2005; 12(46):227-34. Persian.
22. Bakhshandehnosrat S, Qaemi E, Behnampoor N, Rezaeyai M. [Determining the etiological agents in vaginal infections in women referring to Dezyani women hospital in Gorgan]. *J Sabzevar Univ Med Sci.* 2003;10(3):58-65. Persian.
23. Habibypour R, Amirkhani A, Matinnia N. Contamination rate of Trichomonas vaginalis in females referring to Taamin Ejetemayi hospitals in Hamedan in 2005. *Zahedan J Res Med Sci.* 2007;8(4):245-51.
24. Hazrati Kh, Mohammadzadeh H, Mostaghim M, Fereiduni J, Mehri E. [A comparative study sensitivity diagnostic of smear and Diamond culture methods for detection of Trichomonas vaginalis and the relationship between infection and clinical findings]. *J Urmia Univ Med Sci.* 2004;15:7-13. Persian.
25. Moshfe AA, Hosseini S. Comparison of clinical and microscopis diagnosis of Trichomoniasis referred to the Yasouj Women Clinic. *Armaghane Danesh.* 2004;9(1):71-82.
26. Akhlaghi L, Falahati M, Jahani Abianeh M, Ourmazdi H, Amini M. [Study on the prevalence of Trichomonas vaginalis and Candida Albicans in women referred to Robat Karim medical center and a comparative evaluation of Löffler and Diluted Carbol Fuchsin Stains for rapid diagnosis of them]. *Razi J Med Sci.* 2005;12(48):75-87. Persian.
27. Sharbat Daran M, Shefaei Sh, Sami H, Haji Ahmadi M, Ramezan Pour R, Mersadi N, et al. Comparison of clinical presentations, wet smear, Papanicolaou smear with Dorset's culture for diagnosis of Trichomonas vaginalis in doubtful women to Trichomoniasis. *J Babol Univ Med Sci.* 2005;7(3):46-9.
28. Abdolali chalechale, Isaac karimi. The prevalence of Trichomonas vaginalis infection among patients that presented to hospitals in the Kermanshah district of Iran in 2006 and 2007. *Turk J Med Sci.* 2010;40:971-5.
29. Jamali R, Zareikar B, Yousefee S, Ghazanchaei A. Comparison of direct microscopic examination and culture methods sensitivity for diagnosis of Trichomonas vaginalis in Tabriz health care centers visitors. *Yafteh.* 2007;8(4):63-8.
30. Etminan S, Bokaei M. Prevalence of trichomoniasis in women referring to health centers in Yazd. *J Knowledge Health.* 2007;2(3):14-20.
31. Gouya MM, Nabai S. [Prevalence of some sexually transmitted infections in a family planning service]. *Razi J Med Sci.* 2007;14(54):143-50. Persian.
32. Bulbul haghghi N, Ebrahimi H, Delvarian-Zade M, Hasani MR. [Evaluate and compare the clinical and laboratory diagnosis of candida vaginitis in women referred to health centers in the Shahrood city]. *J Shahrekord Univ Med Sci.* 2009;11(3):17-22. Persian.
33. Ziaei Hezarjaribi H, Dalimi A, Ghasemi M, Ghafari R, Esmaili S, Armat S, et al. [Prevalence of comm on sexually transmitted diseases among women referring for Pap smear in Sari, Iran]. *J Mazand Univ Med Sci.* 2013;23(1):19-24. Persian.
34. Rezaeian M, Vatanshenassan M, Rezaie S, Mohebbali M, Niromand N, Niyyati M, et al. Prevalence of Trichomonas vaginalis using parasitological methods in Tehran. *Iran J Parasitol.* 2009;4(4):43-7.
35. Badparva E, Papi OA, kheirandish F, Pornia Y, Azizi M. Sensitivity assessment of direct method for diagnosis of Trichomonas vaginalis in comparison with Dorset culture media. *Yafteh.* 2010;12(1): 25-30.

36. Makvandi S, Zargar Shoushtari Sh. The relationship of cervicovaginal infections in Pap smear samples with some factors in Ahvaz, Iran; an epidemiological study. *Jundishapur J Chronic Dis Care*. 2011;1(1):55-61.
37. Baghchesaraie H, Salmani R, Amini B. Prevalence of *Trichomonas vaginalis* infection among women referred to laboratories in Zanjan, 2010. *J Res Develop in Nurs Midwifery*. 2012;9(1):69-75.
38. Nourian AA, Shabani N, Mousavinasab S, Rahmampour H. Association of *Trichomonas vaginalis* with low birth weight. *J Zanjan Univ Med Sci*. 2011;19(76):84-93.
39. Matini M, Rezaie S, Mohebbali M, Maghsood AH, Rabiee S, Fallah M, et al. Genetic identification of *Trichomonas vaginalis* by using the actin gene and molecular based methods. *Iran J Parasitol*. 2014;9(3):329-35.
40. Talari S, Kazemi B, Hooshyar H, Kazemi F, Arbab M, Talari MR, et al. Detection of drug resistance gene in *Trichomonas vaginalis* by PCR. *Feyz*. 2011;15(1):44-9.
41. Nazari N, Zangeneh M, Moradi F, Bozorgomid A. Prevalence of Trichomoniasis among women in Kermanshah, Iran. *Iran Red Crescent Med J*. 2015;17(3):e23617.
42. Arbabi M, Fakhrieh Z, Delavari M, Abdoli A. Prevalence of *Trichomonas vaginalis* infection in Kashi city, Iran (2012-2013). *Iran J Reprod Med*. 2014;12(7):507-12.
43. Maghsoudi R, Danesh A, Kabiri N, Setorki M, Doudi M. Prevalence of the genital tract bacterial infections after vaginal reconstructive surgery. *Pak J Biol Sci*. 2014;17(9):1058-63.
44. Safayi delouyi Z, Valadkhani Z, Sohrabi M. Analysis the prevalence of *Trichomonas vaginalis* in women clinics of Tehran city's referents by PCR. *Horizon Med Sci*. 2015;20(4):223-9.
45. Habibi A, Nateghi Rostami M, Douraghi M, Dolati M, Hossein Rashidi B, Ahangari R. Frequency of genital infection with *Trichomonas vaginalis* in women referred to gynecology hospital of the city of Qom. *J Dermatol Cosmet*. 2015;6(4):190-219.
46. Van der Pol B. *Trichomonas vaginalis* infection: the most prevalent nonviral sexually transmitted infection receives the least public health attention. *Clin Infect Dis*. 2007;44(1):23-5.
47. Van Der Pol B, Kwok C, Pierre-Louis B, Rinaldi A, Salata RA, Chen PL, et al. *Trichomonas vaginalis* infection and HIV acquisition in African women. *J Infect Dis*. 2008;197(4):548-54.
48. Harp DF, Chowdhury I. Trichomoniasis: evaluation to execution. *Eur J Obstet Gynecol Reprod Biol*. 2011;157(1):3-9.
49. N Poole D, Mc Clelland RS. Global epidemiology of *Trichomonas vaginalis*. *Sex Transm Infect*. 2013;89(6):418-22.
50. Seña AC, Miller WC, Hobbs MM, Schwebke JR, Leone PA, Swygard H, et al. *Trichomonas vaginalis* infection in male sexual partners: implications for diagnosis, treatment, and prevention. *Clin Infect Dis*. 2007;44(1):13-22.
51. Van Der Pol B, Kraft CS, Williams JA. Use of an adaptation of a commercially available PCR assay aimed at diagnosis of chlamydia and gonorrhea to detect *Trichomonas vaginalis* in urogenital specimens. *J Clin Microbiol*. 2006;44(2):366-73.
52. Akbari Z, Matini M. The study of Trichomoniasis in pregnant women attending Hamadan city health centers in 2015. *Avicenna J Clin Microb Infec*. 2017;4(2):e41533.
53. Matini M, Rezaei H, Fallah M, Maghsood AH, Saidijam M, Shamsi-ehsan T. Genotyping, drug susceptibility and prevalence survey of *Trichomonas vaginalis* among women attending gynecology clinics in Hamadan, western Iran, in 2014–2015. *Iran J Parasitol*. 2017;12(1):29-37.
54. Nazari N, Nomani H, Mikaeili A, Hamzavi Y, Mehdaraghi MT, Foroughinia S. The prevalence of *Trichomonas vaginalis* in pap smear samples of women presented to Imam Reza hospital, Kermanshah, Iran from 2006-2012. *Res J Med Sci*. 2016;10(6):653-8.
55. Ahady MT, Safavi N, Jafari A, Mohamadi Z, Abed S, Poursagar S. Prevalence of Trichomoniasis among 18-48 year-old women in northwest of Iran. *Iran J Parasitol*. 2016;11(4):580-4.