



Dietary Fatty Acid Intakes and the Outcomes of Assisted Reproductive Technique in Infertile Women

Maryam Jahangirifar¹, Mahboube Taebi^{2*}, Mohammad Hossein Nasr-Esfahani³, Motahar Heidari-Beni⁴, Gholam Hossein Asgari⁵

1- School of Nursing and Midwifery, Faculty of Medicine, Nursing and Health Sciences, Monash University, Melbourne, Australia

2- Department of Midwifery and Reproductive Health, Isfahan University of Medical Sciences, Isfahan, Iran

3- Department of Animal Biotechnology, Reproductive Biomedicine Research Center, Royan Institute for Biotechnology, ACECR, Isfahan, Iran

4- Child Growth and Development Research Center, Research Institute for Primordial Prevention of Non-Communicable Disease, Isfahan University of Medical Sciences, Isfahan, Iran

5- Department of Community Nutrition, Isfahan University of Medical Sciences, Isfahan, Iran

Abstract

Background: The purpose of this study was evaluating the relationship between fatty acid (FA) intakes and the Assisted Reproductive Technique (ART) outcomes in infertile women.

Methods: In this descriptive longitudinal study, a validated food frequency questionnaire (FFQ) was used to measure dietary intakes among 217 women with primary infertility seeking ART treatments at Isfahan Fertility and Infertility Center, Isfahan, Iran. The average number of total and metaphase II (MII) oocytes, the fertilization rate, the ratio of good and bad quality embryo and biochemical and clinical pregnancy were assessed. Analyses were performed using mean, standard deviation, Chi-square test, ANOVA, ANCOVA, logistic regression.

Results: A total of 140 women were finally included in the study. There was a positive relationship between the average number of total and MII oocytes and the amount of total fatty acids (TFAs), saturated fatty acids (SFAs), monounsaturated fatty acids (MUFAs), polyunsaturated fatty acids (PUFAs), linoleic acids, linolenic acids, and oleic acids intakes, while eicosapentaenoic acids (EPAs) and docosahexaenoic acids (DHAs) intakes had an inverse relationship. Consuming more amounts of TFAs, SFAs, PUFAs, MUFAs, linoleic acids, and oleic acids was associated with the lower fertilization rate, whereas the consumption of linolenic acids and EPAs increased the fertilization rate. The ratio of good quality embryo was directly affected by the amount of PUFAs intakes. Additionally, there was a negative correlation between the amount of SFAs intakes and the number of pregnant women.

Conclusion: TFAs, SFA, PUFA, and MUFA intakes could have both beneficial and adverse impacts on ART outcomes.

Keywords: Assisted reproductive technique, Dietary fats, In vitro fertilization, Infertility, Nutrition assessment.

To cite this article: Jahangirifar M, Taebi M, Nasr-Esfahani MH, Heidari-Beni M, Asgari GhH. Dietary Fatty Acid Intakes and the Outcomes of Assisted Reproductive Technique in Infertile Women. *J Reprod Infertil.* 2021;22(3):173-183. <http://dx.doi.org/10.18502/jri.v22i3.6718>.

Introduction

Infertility is one of the most common challenges faced by reproductive age women today, affecting 10-15% of couples (1). It is

defined as a failure of a couple to conceive after having regular, unprotected sexual intercourse for one year (2). In a recent systematic review by

* Corresponding Author:
Mahboube Taebi,
Department of Midwifery
and Reproductive Health,
Isfahan University of
Medical Sciences, Isfahan,
Iran
E-mail:
m_taebi@nm.mui.ac.ir

Received: Mar. 25, 2021

Accepted: July. 02, 2021

Dirkavand-Moghaddam et al., the overall rate of infertility in Iran was reported to be 13.2% (3). Although using ART such as in vitro fertilization (IVF) could be of great benefit, only 37.1% of fresh, nondonor ART cycles in women under 35 referred to the United States clinics in 2014 have resulted in live births (4), which shows there might be some unknown factors to be discovered.

Improper nutrition is known as a growing problem all over the world and according to estimates, by 2030, up to 57.8% of the world's adult population (3.3 billion people) could be either overweight or obese (5) which emphasizes the necessity for focusing on this area of research. Furthermore, recent studies demonstrated that maternal nutritional status has profound effects on reproductive health (6) and the metabolic changes could be reflected in the follicular fluid of the dominant follicles (7). In fact, oocyte quality, which is one of the determinants of successful embryonic development, is dependent on the ovarian and follicular environment changes, especially those stressors induced by changes in nutrients (8, 9).

FAs including SFA, MUFA, and PUFA perform a crucial role in oocyte maturation and embryo development (10). It was shown that linoleic acid (A type of PUFA) could enhance oocyte competence and embryo development beyond 4-cell stage (11); however, some studies demonstrated that high levels of LIFA could negatively influence oocyte and embryo quality (12-14). Regarding the effect of oleic acid (A type of MUFA) on intracytoplasmic sperm injection (ICSI) outcomes, majority of studies showed that this FA intake could contribute to normal oocyte and embryo development, notwithstanding some conflicting reports (10, 15-17). Moreover, there are a number of other studies with controversial results about the effects of other types of FA intakes on reproductive health which require further investigation (13, 18). In addition, despite conducting many studies on the effects of FA intakes on oocyte competence and embryo development in animals, there is a distinct lack of research on human reproductive response to FA intakes. Therefore, this study was designed to evaluate the relationship between dietary FA intakes and the outcome of ART in infertile women.

Methods

Participants: This study was a descriptive longi-

tudinal study that was performed at Isfahan Fertility and Infertility Center, Isfahan, Iran. A simple sampling design was used and 217 infertile women seeking treatments including IVF and ICSI were enrolled into the study from August 2015 to January 2016. The inclusion criteria were having primary female infertility (Unexplained infertility or ovarian failure), no significant change in diet during the last 3 months or a special diet, not having diseases affecting metabolism like diabetes, hypothyroidism, *etc.*, not using drugs affecting the metabolism of macro and micro nutrients including blood sugar-lowering and lipid-lowering drugs, the absence of anatomic abnormalities, endometriosis and surgery in the uterus and tubes, not using a surrogate, not using alcohol, and non-smoking. Based on the findings of a previous study, the sample size for each tertile was calculated to be 65 for $d=0.6P$ and $p=40\%$, and considering 10% participant attrition, the required total sample size was approximately 215 (19).

Dietary assessment: A validated food frequency questionnaire (FFQ) including 168 food items was used to measure dietary intakes (20, 21). When participants came to choose their treatment plans, they were asked to report how often and how much they had consumed each of the foods during the previous year. Then, amount of food items was converted into grams and calculated for one day and entered into Nutritionist IV software for each person. This software calculated all food ingredients such as FAs. Next, the amount of different types of FAs for each person was calculated on the basis of food source by Excel (Microsoft office 2016) which was entered into SPSS. Also, the daily physical activity as a confounding variable was assessed by using the short form of International Physical Activity Questionnaire (IPAQ). The amount of activity was computed by weighting each type of activity by its energy requirements defined in metabolic equivalent of task (MET) *min/day*. Then, it was converted into (MET) *h/week* and reported. These questionnaires have been validated in previous studies (20, 22-24). BMI was calculated as weight in kg divided by the square of height in meters (kg/m^2). The waist circumference was measured with a non-stretch tape to the nearest 0.5 *cm* between the lowest rib margin and the iliac crest wearing minimal clothing (25).

Ovulation induction: In the present study, patients followed the suppression protocol. On the second

day of last menstrual period, recombinant follicle stimulating hormone (Gonal-F) (Serono, Switzerland) in combination with human menopausal gonadotropin (Menogon) (Ferring Pharmaceuticals, Germany) was commenced, when transvaginal sonography showed absence of ovarian cysts. When the size of dominant follicles reached 17-18 mm, ovulation was induced with 10,000 IU hCG. Transvaginal oocyte retrieval was done 36 hr later.

Laboratory assessment: In the laboratory, MII oocytes were recorded. Fertilization rate was defined as the ratio of zygotes with two pronuclei observed 18 hr after insemination divided by the number of oocytes inseminated. Embryos were scored by using a four-point score on day 3 (good quality=score 4, bad quality=score 2). All cleaved embryos were assigned 1 point, and an additional point was added for each of the following features: absence of fragmentation (or fragmentation involving 25% of embryonic surface), absence of irregularities in blastomere size or shape, 8-cell stage on Day 3. Biochemical pregnancy was defined as the presence of β hCG in serum 12 days after embryo transfer. Clinical pregnancy was confirmed by detection of one or more gestational sacs during transvaginal scan 3 weeks after embryo transfer.

Statistical analysis: Collected data were analyzed using SPSS software version 20.0 (IBM Corp, New York). For demographic characteristics, ANOVA and Chi-square tests were used to assess differences in continuous and categorical variables, respectively. ANOVA and ANCOVA analyses were used to determine the association between tertiles of different types of FA intakes and fertility markers including the average number of total oocytes, the average number of MII oocytes and the ratio of good and poor quality embryo. The model was adjusted for age, marriage age, BMI, waist circumference, supplement consumption, Metformin consumption, and physical activity. Logistic regression analysis was used (With covariates referred above) to calculate adjusted odds ratios (AOR) and 95% CI for assessing the association between different types of FA intakes and biochemical and clinical pregnancy. A p-value of less than 0.05 was considered statistically significant.

This study was conducted according to the guidelines laid down in the Declaration of Helsinki and all procedures involving research study participants were approved by the Ethics Commit-

tee of Isfahan University of Medical Sciences (Ethical code: IR.MUI.Rec.1394.3.475). Written informed consent was obtained from all subjects.

Results

Of the 217 infertile women participated in this study, 77 cases were excluded (Owing to spontaneous pregnancy, cancellation of treatment cycle due to poor or excessive response to stimulation, ovulation before oocyte retrieval, diagnosis of male factor infertility at later stages, refusal to cooperate during the study period and not completing the follow-up). Figure 1 and table 1 show the details of sample recruitment and baseline characteristics, respectively. A total of 140 women were finally included in the final analysis. The majority of women with lower SFA intake who were categorized into the first and third tertiles of linoleic acid intake were housewives ($p=0.005$, $p=0.032$, respectively). Moreover, women who had higher education were more likely to have a diet rich in linolenic acid ($p=0.043$).

In table 2, the results showed there was a significant increase in the average number of total and MII oocytes in the higher tertile of TFA, SFA, MUFA, PUFA, linoleic acid, and linolenic acid intakes after adjusting for confounders. The amount of oleic acid intake had a direct relationship with the average number of total and MII oocytes before ($p=0.006$, $p=0.000$, respectively) and after ($p=0.007$, $p=0.000$, respectively) adjusting for confounders. It was shown that women in the higher tertile of EPA and DHA had lower number of total and MII oocytes after controlling for confounding variables ($p=0.001$, $p=0.001$, respectively).

Furthermore, our data showed (After adjusting for confounders) consuming more amounts of dietary TFA, SFA, PUFA, MUFA, linoleic acid, and oleic acid was associated with the lower fertilization rate, whereas there was a positive association between the consumption of linolenic acid and EPA and the fertilization rate ($p=0.007$, $p=0.003$). The ratio of good quality embryo was directly associated by the amount of dietary PUFA, which was not statically significant after adjusting for the confounders. Additionally, there was a negative correlation between the amount of SFA intake and the number of pregnant women (Biochemical and clinical pregnancy, OR=0.06, CI=0.007-0.572, $p=0.001$) which remained significant after considering confounders (OR=0.008, CI=0.00-1.49, $p=0.002$).




Figure 1. Participant recruitment

Table 1. Demographic features of infertile women by tertiles

Variables	TFA				SFA				MUFA			
	T1	T2	T3	p ^d	T1	T2	T3	p	T1	T2	T3	p
Age, year (M±SD) ^a	32.11±5.67	32.87±4.95	32.23±5.01	0.75	31.26±5.82	32.98±5.23	32.95±4.35	0.189	33.06±5.57	32.42±4.95	31.74±5.06	0.475
Age at marriage, year (M±SD)	23.59±5.84	24±5.40	23.81±6.34	0.944	23.15±5.73	23.87±5.65	24.36±6.16	0.607	24.82±6.21	23.63±5.70	22.96±5.55	0.298
BMI, weight (kg ^b)/ height (m ²) (M±SD)	28.73±5.32	27.74±4.69	27.70±4.68	0.528	28.99±5.27	26.91±4.48	28.23±4.74	0.119	28.21±4.87	28.04±5.34	27.89±4.51	0.952
Waist circumference, cm (M±SD)	83.65±10.48	83.47±11.09	83.06±9.73	0.962	84.46±10.74	81.34±10.75	84.41±9.51	0.251	83.33±9.33	84.08±12.78	82.77±8.71	0.828
Hip circumference, cm (M±SD)	105.04±16.76	104.36±8.70	105.59±9	0.884	104.83±16.56	104.70±7.37	105.47±10.46	0.947	105.76±9.58	103.19±16.66	106.06±7.73	0.445
Infertility duration, year (M±SD)	6.78±4.89	6.89±4.17	6.95±4.81	0.985	6.59±4.58	7.32±5.23	6.69±3.95	0.707	6.80±4.48	7.17±4.70	6.64±4.68	0.855
Physical activity, met-h/week, (M±SD)	6.11±8	15.06±21.43	14.90±33.73	0.117	5.61±7.65	15.16±21.43	15.29±33.68	0.080	6.57±8.62	11.98±14.57	17.53±36.97	0.084
Education (%)												
Below diploma	14(30.4%)	13(27.7%)	9(19.1%)		14(30.4%)	12(25.5%)	10(21.3%)		11(23.9%)	15(31.9%)	10(21.3%)	
Diploma	13(28.3%)	20(42.6%)	12(25.5%)	0.119	13(28.3%)	20(42.6%)	12(25.5%)	0.220	13(28.3%)	20(42.6%)	12(25.5%)	0.080
Academic	19(41.3%)	14(29.8%)	26(55.3%)		19(41.3%)	15(31.9%)	25(53.2%)		22(47.8%)	12(25.5%)	25(53.2%)	
The cause of infertility (%)												
Ovarian	38(82.6%)	32(68.1%)	35(74.5%)	0.269	37(80.4%)	35(74.5%)	33(70.2%)	0.520	34(73.9%)	33(70.2%)	38(80.9%)	0.482
Idiopathic	8(17.4%)	15(31.9%)	12(25.5%)		9(19.6%)	12(25.5%)	14(29.8%)		12(26.1%)	14(29.8%)	9(19.1%)	
Employment status (%)												
Housewife	40(87%)	37(78.7%)	33(70.2%)	0.144	42(91.3%)	38(80.9%)	30(63.8%)	0.005	37(80.4%)	39(83%)	34(72.3%)	0.423
Employed	6(13%)	10(21.3%)	14(29.8%)		4(8.7%)	9(19.1%)	17(36.2%)		9(19.6%)	8(17%)	13(27.7%)	

Table 1. Demographic features of infertile women by tertiles

Variables	PUFA				Oleic acids				Linoleic acids			
	T1	T2	T3	P	T1	T2	T3	p	T1	T2	T3	p
Age, year (M±SD ^a)	32.80±5.06	31.94±5.27	32.49±5.31	0.719	32.80±5.80	32.59±4.49	31.83±5.28	0.637	32.93±5.31	32.06±5.03	32.23±5.29	0.696
Age at marriage, year (M±SD)	24.02±5.61	23.74±5.51	23.63±6.44	0.949	24.73±6.36	23.64±5.86	23.04±5.24	0.367	23.78±6.12	24.28±5.18	23.34±6.24	0.742
BMI, weight (kg ^b)/ height (m ^c) ² (M±SD)	27.99±4.33	27.57±5.32	28.56±4.99	0.618	28.42±5.41	28.23±4.70	27.49±4.57	0.626	27.88±4.99	28.44±5.41	27.81±4.27	0.796
Waist circumference, cm (M±SD)	83.26±9.13	83.32±12.15	83.59±9.81	0.987	83.81±10.80	84.02±10.81	82.36±9.65	0.703	83.33±10.24	84.94±11.66	81.91±9.07	0.369
Hip circumference, cm (M±SD)	102.83±16.34	104.57±9.33	107.55±8.51	0.156	107.13±10.53	102.62±15.33	105.30±8.86	0.188	102.48±16.60	106.02±9.63	106.45±7.78	0.216
Infertility duration, year (M±SD)	7.16±4.86	6.24±4.05	7.22±4.88	0.517	6.88±4.42	6.51±4.16	7.23±5.21	0.750	7.54±4.74	5.96±4.27	7.14±4.72	0.225
Physical activity, met-h/week, (M±SD)	7.57±9.90	11.60±15.76	16.94±36.39	0.163	8.39±9.61	9.89±13.56	17.84±37.23	0.119	8.42±10.70	10.64±15.53	17.06±36.32	0.192
Education (%)												
Below diploma	15(32.6%)	11(23.4%)	10(21.3%)		14(30.4%)	14(29.8%)	8(17%)		15(32.6%)	10(21.3%)	11(23.4%)	
Diploma	16(34.8%)	11(23.4%)	18(38.3%)	0.243	13(28.3%)	14(29.8%)	18(38.3%)	0.550	15(32.6%)	13(27.7%)	17(36.2%)	0.500
Academic	15(32.6%)	25(53.2%)	19(40.4%)		19(41.3%)	19(40.4%)	21(44.7%)		16(34.8%)	24(51.1%)	19(40.4%)	
The cause of infertility (%)												
Ovarian	35(76.1%)	32(68.1%)	38(80.9%)	0.352	34(73.9%)	33(70.2%)	38(80.9%)	0.482	35(76.1%)	34(72.3%)	36(76.6%)	0.874
Idiopathic	11(23.9%)	15(31.9%)	9(19.1%)		12(26.1%)	14(29.8%)	9(19.1%)		11(23.9%)	13(27.7%)	11(23.4%)	
Employment status (%)												
Housewife	40(87%)	32(68.1%)	38(80.9%)	0.077	39(84.8%)	36(76.6%)	35(74.5%)	0.442	40(87%)	31(66%)	39(83%)	0.32
Employed	6(13%)	15(31.9%)	9(19.1%)		7(15.2%)	11(23.4%)	12(25.5%)		6(13%)	16(34%)	8(17%)	

Table 1. Demographic features of infertile women by tertiles

Variables	Linolenic acids				EPA				DHA			
	T1	T2	T3	p	T1	T2	T3	p	T1	T2	T3	p
Age, year (M±SD ^a)	33.18±5.48	31.85±4.44	32.62±5.68	0.478	-	32.43±5.21	31.75±4.99	0.798	-	32.36±5.21	33.50±6.09	0.600
Age at marriage, year (M±SD)	24.54±6.28	23.22±4.55	24.11±6.52	0.546	-	23.75±5.82	25.50±7.14	0.556	-	23.63±5.77	27.50±6.71	0.113
BMI, weight (kg ^b)/ height (m ^c) ² (M±SD)	29.19±4.98	28.14±4.82	26.65±4.70	0.051	-	28.07±4.93	27.09±2.41	0.735	-	28.06±4.96	27.61±2.36	0.841
Waist circumference, cm (M±SD)	85.20±9.34	83.29±9.77	81.38±11.82	0.224	-	83.38±10.47	84±7.39	0.907	-	83.30±10.53	85.50±6.25	0.614
Hip circumference, cm (M±SD)	104.09±16.64	106±9.92	104.20±8.26	0.701	-	104.94±12.11	107±3.82	0.736	-	104.92±12.17	106.83±5.60	0.703
Infertility duration, year (M±SD)	7.26±4.60	6.72±4.42	6.60±4.62	0.775	-	6.94±4.64	4.62±2.29	0.323	-	6.98±4.66	4.41±1.91	0.182
Physical activity, met-h/week, (M±SD)	6.79±9.77	17.14±35.37	12.13±18.62	0.126	-	12.20±24.10	7.50±9	0.699	-	11.86±24.11	16.65±16.25	0.633
Education (%)												
Below diploma	17(38.6%)	8(17.4%)	8(17.8%)		-	36(26.5%)	0(0%)		-	36(26.9%)	0(0%)	
Diploma	15(34.1%)	13(28.3%)	16(35.6%)	0.043	-	44(32.4%)	1(25%)	0.336	-	44(32.8%)	1(16.7%)	0.100
Academic	12(27.3%)	25(54.3%)	21(46.7%)		-	56(41.2%)	3(75%)		-	54(40.3%)	5(83.3%)	
The cause of infertility (%)												
Ovarian	34(77.3%)	36(78.3%)	34(75.6%)	0.953	-	102(75%)	3(75%)	1	-	100(74.6%)	5(83.3%)	0.630
Idiopathic	10(22.7%)	10(21.7%)	11(24.4%)		-	34(25%)	1(25%)		-	34(25.4%)	1(16.7%)	
Employment status (%)												
Housewife	37(84.1%)	35(76.1%)	34(75.6%)	0.547	-	106(77.9%)	4(100%)	0.289	-	106(79.1%)	4(66.7%)	0.468
Employed	7(15.9%)	11(23.9%)	11(24.4%)		-	30(22.1%)	0(0%)		-	28(20.9%)	2(33.3%)	

a: Standard deviation, b: Kilogram, c: Meter, d: p-value from one-way analysis of variance for continuous quantitative variables and from Chi-square test for categorical variables

The ORs and CIs for the biochemical and clinical pregnancy in the third tertile of EPA and DHA were not reportable and so deliberately omitted from table 2.

Discussion

The aim of the present study was to evaluate the relationship between dietary fatty acid intakes and the outcome of ART in infertile women. The re-

sults showed that consumption of different types of FAs could be related with ART outcomes in different ways.

Leroy et al. were one of the first who conducted a study with the aim of examining the effect of metabolic changes on the follicular microenvironment and potential consequences for oocyte and embryo quality. They carried out a study on high-producing dairy cows in negative energy balance

Table 2. Comparison of the ART outcomes by tertiles of FA intakes in infertile women

Variables	The average number of total oocytes		The average number of MII oocyte		The fertilization rate		The ratio of good quality embryos		The ratio of poor quality embryos		Biochemical pregnancy [OR(CI)] ^e		Clinical pregnancy [OR(CI)]	
	Crude (M±SD) ^b	Adjust ^a (M±SE) ^c	Crude (M±SD)	Adjust (M±SE)	Crude (M±SD)	Adjust (M±SE)	Crude (M±SD)	Adjust (M±SE)	Crude (M±SD)	Adjust (M±SE)	Crude	Adjust	Crude	Adjust
TFA														
T ₁	7.80±5.5	8.20±1.3	6.93±5.3	7.03±1.1	0.67±0.3	0.69±0.05	0.18±0.3	0.19±0.05	0.26±0.3	0.28±0.05	1	1	1	1
T ₂	10.76±8.7	10.78±1.1	8.68±6.9	8.73±1	0.70±0.3	0.69±0.04	0.24±0.3	0.25±0.04	0.30±0.3	0.29±0.05	0.42 (0.12-1.49)	0.50 (0.10-2.33)	0.42 (0.12-1.49)	0.50 (0.10-2.33)
T ₃	11.34±9.6	10.92±1.3	9.74±8.2	9.58±1.1	0.68±0.2	0.65±0.05	0.16±0.2	0.13±0.05	0.38±0.4	0.37±0.05	0.44 (0.10-1.84)	0.20 (0.02-1.89)	0.44 (0.10-1.84)	0.20 (0.02-1.89)
p ^d	0.085	0.001	0.148	0.001	0.912	0.007	0.363	0.358	0.23	0.635	0.327	0.370	0.327	0.370
SFA														
T ₁	8.10±6.2	7.92±1.2	7.13±5.2	6.75±1	0.73±0.3	0.74±0.05	0.27±0.3	0.28±0.04	0.31±0.3	0.33±0.05	1	1	1	1
T ₂	10.93±8.9	11.13±1.2	8.95±7.4	9.23±1	0.65±0.3	0.65±0.05	0.18±0.2	0.19±0.04	0.26±0.3	0.25±0.05	0.02 (0.003-0.254)	0.01 (0.001-0.18)	0.02 (0.003-0.25)	0.016 (0.001-0.18)
T ₃	10.87±9.1	10.84±1.2	9.27±7.9	9.37±1	0.66±0.3	0.65±0.05	0.13±0.2	0.11±0.04	0.38±0.4	0.36±0.05	0.06 (0.007-0.572)	0.008 (0.00-1.49)	0.06 (0.007-0.57)	0.008 (0.00-1.49)
p	0.17	0.001	0.282	0.001	0.417	0.003	0.056	0.064	0.193	0.44	0.001	0.002	0.001	0.002
MUFA														
T ₁	8.93±7.8	9.99±1.3	7.63±6.7	8.40±1	0.66±0.3	0.68±0.05	0.14±0.2	0.14±0.05	0.25±0.3	0.26±0.05	1	1	1	1
T ₂	9.38±7.6	9.45±1.2	7.97±6.2	8.01±1	0.72±0.3	0.73±0.04	0.20±0.3	0.20±0.04	0.35±0.3	0.35±0.05	1.80 (0.46-6.97)	1.70 (0.34-8.52)	1.80 (0.46-6.97)	1.70 (0.34-8.52)
T ₃	11.61±9.1	10.51±1.3	9.76±7.9	8.96±1.1	0.66±0.2	0.63±0.05	0.24±0.3	0.24±0.05	0.34±0.3	0.33±0.05	1.74 (0.43-6.97)	2.07 (0.32-13.14)	1.74 (0.43-6.97)	2.07 (0.32-13.14)
p	0.243	0.003	0.286	0.004	0.533	0.003	0.251	0.542	0.281	0.625	0.652	0.721	0.652	0.721
PUFA														
T ₁	8.69±7.4	9.21±1.2	7.52±6.6	7.85±1	0.66±0.3	0.67±0.05	0.12±0.2	0.11±0.04	0.28±0.3	0.30±0.05	1	1	1	1
T ₂	10.85±8.3	10.80±1.2	8.95±6.6	8.90±1	0.75±0.3	0.65±0.04	0.19±0.3	0.19±0.04	0.37±0.3	0.38±0.05	1.52 (0.39-5.91)	2.21 (0.39-12.42)	1.52 (0.39-5.91)	2.21 (0.39-12.42)
T ₃	10.38±9	9.92±1.2	8.89±7.7	8.62±1.1	0.63±0.3	0.61±0.05	0.27±0.3	0.27±0.04	0.29±0.3	0.27±0.05	1.60 (0.39-6.45)	1.87 (0.26-13.48)	1.60 (0.39-6.45)	1.87 (0.26-13.48)
p	0.419	0.003	0.539	0.003	0.118	0.001	0.044	0.125	0.333	0.45	0.775	0.658	0.775	0.658
Oleic acids														
T ₁	6.95±5.8	7.24±1.2	5.84±5.3	6.03±1	0.68±0.4	0.69±0.05	0.17±0.3	0.18±0.05	0.25±0.3	0.26±0.05	1	1	1	1
T ₂	10.70±7.5	11±1.1	9.61±7.3	9.84±1	0.66±0.3	0.68±0.04	0.14±0.2	0.14±0.04	0.39±0.3	0.40±0.05	1.16 (0.30-4.43)	1.81 (0.35-9.32)	1.16 (0.30-4.43)	1.81 (0.35-9.32)
T ₃	12.23±10	11.64±1.2	9.87±7.4	9.46±1	0.70±0.2	0.67±0.05	0.26±0.3	0.26±0.05	0.30±0.3	0.27±0.05	2.16 (0.56-8.25)	2.68 (0.40-17.80)	2.16 (0.56-8.25)	2.68 (0.40-17.80)
p	0.006	0.000	0.007	0.000	0.885	0.008	0.117	0.245	0.119	0.158	0.480	0.589	0.480	0.589
Linoleic acids														
T ₁	8.76±6.9	9.27±1.2	7.50±6	7.88±1	0.74±0.2	0.67±0.05	0.17±0.2	0.17±0.04	0.30±0.3	0.32±0.05	1	1	1	1
T ₂	10.87±9.4	11.16±1.2	9.25±8	9.43±1	0.68±0.3	0.65±0.04	0.19±0.3	0.19±0.04	0.31±0.3	0.31±0.05	1.48 (0.40-5.49)	1.96 (0.40-9.60)	1.48 (0.40-5.49)	1.96 (0.40-9.60)
T ₃	10.29±8.2	9.50±1.3	8.61±6.8	8.06±1.1	0.62±0.3	0.61±0.05	0.22±0.3	0.22±0.05	0.33±0.3	0.32±0.05	1.08 (0.29-4.01)	0.64 (0.10-3.84)	1.08 (0.29-4.01)	0.64 (0.10-3.84)
p	0.446	0.002	0.475	0.002	0.405	0.002	0.632	0.906	0.907	0.979	0.822	0.407	0.822	0.407
Linolenic acids														
T ₁	8.44±8.2	9.41±1.2	7.22±6.9	7.87±1.1	0.63±0.4	0.64±0.05	0.26±0.2	0.27±0.04	0.24±0.3	0.25±0.05	1	1	1	1
T ₂	11.24±9.7	10.81±1.2	9.24±7.9	8.92±1	0.71±0.3	0.70±0.05	0.18±0.3	0.18±0.04	0.29±0.3	0.38±0.05	1.14 (0.29-4.36)	0.8 (0.16-4.18)	1.14 (0.29-4.36)	0.81 (0.16-4.18)
T ₃	10.35±6.8	9.81±1.2	9.02±6.2	8.69±1	0.71±0.2	0.70±0.05	0.15±0.3	0.15±0.04	0.39±0.3	0.28±0.05	0.28 (0.06-1.17)	0.29 (0.05-1.50)	0.28 (0.06-1.17)	0.29 (0.05-1.50)
p	0.269	0.005	0.335	0.005	0.393	0.007	0.148	0.265	0.09	0.302	0.137	0.307	0.137	0.307
EPA														
T ₁	-	-	-	-	-	-	-	-	-	-	-	-	-	-
T ₂	10.11±8.3	10.13±0.7	8.55±7	8.57±0.6	0.68±0.3	0.68±0.03	0.20±0.3	0.20±0.02	0.31±0.3	0.31±0.03	1	1	1	1
T ₃	5.75±2.2	5.03±4	5.25±2.5	4.70±3.4	0.71±0.2	0.71±0.1	0.05±0.1	0.03±0.1	0.33±0.5	0.30±0.2	-	-	-	-
p	0.299	0.001	0.352	0.001	0.84	0.003	0.298	0.539	0.929	0.913	0.999	0.999	0.999	0.999
DHA														
T ₁	-	-	-	-	-	-	-	-	-	-	-	-	-	-
T ₂	10.08±8.4	10.08±0.7	8.52±7.1	8.5±0.3	0.68±0.3	0.68±0.03	0.20±0.3	0.20±0.02	0.31±0.3	0.31±0.03	1	1	1	1
T ₃	7.66±3.5	7.77±3.3	7±3.7	7.36±0.4	0.78±0.2	0.78±0.1	0.03±0.1	0.03±0.1	0.44±0.4	0.39±0.1	-	-	-	-
p	0.483	0.001	0.602	0.001	0.403	0.002	0.152	0.373	0.353	0.781	0.999	0.999	0.999	0.999

a: Adjusted for age, age at marriage, BMI, waist circumference, physical activity, total energy intake, supplement consumption, duration of metformin consumption. b: Standard deviation. c: Standard error. d: p trends from ANOVA analysis for crude and from ANCOVA analysis for adjust in quantitative variables and p trends from logistic regression analysis for qualitative variables. e: OR (CI): odds ratio and 95% interval confidence calculated by logistic regression analysis

in early post-partum and it was shown that the metabolic changes could be reflected in the follicular fluid of the dominant follicles (7). Since there is much similarity between bovine and human reproductive physiology (26), it could be concluded that dietary FA intakes in humans could be associated with reproductive function by changing ovarian activity, follicular growth, corpus luteum function, and the uterine environment (27).

It was revealed that dietary oleic acids could have a favorable impact on the number of re-

trieved oocytes. In the previous study by Salehi et al., it was shown that oleic acids through reducing the release of gonadotropin releasing hormone induced luteinizing hormone in dairy cows (Not affecting basal LH) could directly modulate gonadotropin release from the pituitary and affect ovulation (28). In another study with the purpose of determining the effect of FAs in serum and follicular fluid on ICSI outcomes, it was shown that the mean number of retrieved oocytes was positively associated with serum levels of oleic acid. Simi-

larly, Bilby et al. demonstrated that there were more collected oocytes from dairy cows fed oleic acid in comparison with cows fed trans oleic acids, linoleic acid, or linolenic acid (29). In spite of its beneficial effect, it was found that more consumption of oleic acid causes the reduction in fertilization rate. Consistent with our results, Jorritsma et al. have demonstrated that the exposure of oocyte to high concentration of oleic acid delayed the progression of meiosis and reduced successive fertilization, cleavage, and blastocyst development rates (17). Nevertheless, there are some data in support of positive effects of oleic acid on fertilization (15) which requires more future studies.

Two types of PUFAs, including linoleic acids and linolenic acids, play an important role in reproductive physiology, affecting oocyte quality (30) and in our study, their amounts in diet have shown a direct association with the number of oocytes. As PUFAs are the essential components of membrane lipids, they increase rapidly with each cell division, as it was shown that there is a 74% increase in membrane surface area in transition from the one to the four cell stage in 2-cell divisions (11, 31). In similar studies conducted on goats and pigs, it was demonstrated that adding 50 μM of LNFA to maturation medium increased meiotic maturation rate (32, 33). Marei et al. showed treatment of cumulus-oocyte complexes with 50 μM of linolenic acid significantly increased the percentage of oocytes at the MII stage compared with untreated controls in dairy cattle, which was attributed to the mitogen-activated protein kinase (MAPK) pathway and indirectly to prostaglandin E2 (PGE2) synthesis (34). In fact, PGE2 by stimulating MAPK1 and MAPK2 phosphorylation in both oocytes and cumulus cells through the elevation of cyclic adenosine monophosphate (cAMP) levels plays a key role in oocyte maturation and cumulus expansion (34, 35). In mice, inhibition of PGE production by using its inhibitors or inactivation of encoding genes of the PGE2 receptor was found to stop cumulus expansion and oocyte maturation and considerably decrease fertilization rates *in vitro* (34), which is in parallel with our results concerning the direct relation of linolenic acid intake with the fertilization rate.

Regarding linoleic acid, it was demonstrated this type of FA can serve as a precursor for two series of prostaglandins (With a possible effect on ovulation) (36) and stimulate protein kinase C (37)

which is critical in cell growth and differentiation (38). Despite its positive role in oocyte competence, our study showed the low fertilization rate in women with a diet high in linoleic acid. In a prospective study conducted on 54 women undergoing ICSI, it was reported that high levels of linoleic acid significantly decrease fertilization rate in ICSI cycles. The mechanism behind this was attributed to the activity of secretory phospholipase A2 (sPLA2), since linoleic acid release from a phospholipid molecule is controlled by phospholipases (39). It was reported that sPLA2 enzyme activity causes a number of changes including inflammation and cell degeneration (40).

In this study, the amount of dietary TFA and SFA positively influenced the number of MII oocytes. However, previous studies have shown that exposure to excessive dietary SFAs and high fat diet could result in oocyte mitochondrial damage, which consequently induces oxidative stress (41) and endoplasmic reticulum (ER) stress (42, 43). Likewise, Leroy et al. reported the addition of SFAs such as stearic and palmitic acids to the follicular fluid of dairy cows during *in vitro* maturation delayed meiosis progression, resulting in a significantly greater number of MI oocytes and a relatively lower number of MII oocytes (44). However, Aardema et al. in their study with the aim of examining the effect of three types of fatty acids on bovine oocyte developmental competence showed that oleic acid compensates for the adverse effects of palmitic and stearic acids (16). Therefore, it could be concluded that the positive effect of SFAs on the number of MII oocytes in this study may be confounded by the other types of fatty acids. Moreover, it was demonstrated that women with more consumption of SFAs and TFAs had lower fertilization rate. It was previously shown that lipotoxicity as a result of high concentration of SFAs in maturation media and high-fat diets by induction of ER stress pathway genes and alteration of mitochondrial membrane potential could decrease the fertilization rate (43, 45). In fact, exposure of the ER to high levels of free fatty acids as the main site for the biosynthesis of steroids, cholesterol and other lipids causes structural changes and failures in performing its functions (46).

Women with a diet high in EPA and DHA have the fewer MII oocytes. Similarly, Hammiche et al. demonstrated high intakes of EPA and DHA reduced E2 response and the number of follicles after ovarian stimulation (47). Correspondingly, a

study on rats fed a diet high in EPA and DHA showed a decline in frequency of ovulation (48). The reduction in PGF2 α involved in follicle growth and ovulation which was attributed to EPA and DHA may relatively explain the reduced number of MII oocytes in this study (49, 50). However, there is some inconsistency in view of its deleterious or salutary effect on follicle development which needs further investigation (51). In spite of the inverse relationship between EPA intake and the number of MII oocytes in this study, it has been revealed that consuming MUFA such as EPA could increase the fertilization rate. Indeed, it was demonstrated the gene expression of insulin-like growth factor-I (IGF-I) in granulosa cells increased by EPA (52) could improve the fertilization rate and embryo development (53).

The present study demonstrated that consumption of PUFAs could improve embryo quality. However, after adjusting for confounders, the effect was not significant. Cerri et al. showed cows fed linoleic acid and trans-octadecenoic acids compared with cows fed palm oil had the greater proportion of excellent-, good-, and fair- quality embryos (54). In another study, it was reported the higher intake of linoleic acid and DHA could improve the embryo morphology in women undergoing IVF/ICSI treatment (55).

The final result obtained from this study is the negative relationship between SFA intake and biochemical and clinical pregnancy. Correspondingly, the result from a study conducted on human embryos indicated embryos developed beyond the 4-cell stage had higher concentrations of the unsaturated fatty acids such as linoleic acid, oleic acid, and lower concentration of SFAs. Therefore, the availability of particular fatty acids *in vivo* could potentially influence IVF success following transfer (11). Also, in another study carried out on lactating dairy cattle, embryos of cows fed diets enriched in unsaturated fatty acids in comparison with SFAs have shown more development (55), which may affect the success of conception.

The strengths of this study include using validated questionnaires, evaluating several reproductive outcomes, and following up participants for 5 weeks. Moreover, limitations of this study are uncontrollability of male epigenetic factors like nutrition, measurement errors in data collected using FFQ such as intake-related bias, person-specific bias, and within-person variation (20), the embryologist's mistake in IVF and ICSI procedures, and the limitations of observational studies such as

confounding factors (56). Although a reasonable number of confounders were considered in this study, there were lots of factors that could not be measured due to the limitation of time and budget.

Conclusion

Our study demonstrated that TFAs, SFA, PUFA, and MUFA intakes could have both beneficial and adverse impacts on ART outcomes. Due to glaring inconsistencies in research into the relationship between dietary fatty acids and reproductive outcomes, there is a need for conducting further research. Moreover, since other food ingredients can affect reproductive outcomes, the relation of dietary nutrient intakes with ART outcomes should be evaluated in future studies.

Acknowledgement

We express our gratitude to Vice-Chancellor for Research of Isfahan University of Medical Sciences and Isfahan Fertility and Infertility Center. We would like to thank for their cooperation as well as the assistance of participants who made this study possible.

Conflict of Interest

Authors declare no conflict of interest.

References

1. Rizk B, Olsen ME. Challenging Office Cases in Obstetrics and Gynecology. Johnson City: Jaypee Brothers Medical Publishers Ltd; 2016. p. 65-65.
2. Ramesh B, Kurkuri SN. Role of combined hystero-laparoscopy in the evaluation of female infertility as one step procedure. *Int J Reprod Contraception Obstet Gynecol.* 2016;5:396-401.
3. Dirkevand Moghaddam A, Delpisheh A, Sayehmiri K. The prevalence of infertility in Iran, a systematic review. *Iran J Obstet Gynecol Infertil.* 2013;16(8):1-7.
4. Centers for Disease Control and Prevention, American Society for Reproductive Medicine, Society for Assisted Reproductive Technology. 2014 Assisted reproductive technology: National summary report. Atlanta (GA): US Dept of Health and Human Services; 2016. Report No.: GS-23F-8144H.
5. Kelly T, Yang W, Chen CS, Reynolds K, He J. Global burden of obesity in 2005 and projections to 2030. *Int J Obes (Lond).* 2008;32(9):1431-7.
6. Mmbaga N, Luk J. The impact of preconceptual diet on the outcome of reproductive treatments. *Curr Opin Obstet Gynecol.* 2012;24(3):127-31.
7. Leroy JL, Vanholder T, Delanghe JR, Opsomer G,

- Van Soom A, Bols PE, et al. Metabolic changes in follicular fluid of the dominant follicle in high-yielding dairy cows early post partum. *Theriogenology*. 2004;62(6):1131-43.
8. Gu L, Liu H, Gu X, Boots C, Moley KH, Wang Q. Metabolic control of oocyte development: linking maternal nutrition and reproductive outcomes. *Cell Mol Life Sci*. 2015;72(2):251-71.
 9. Pohlmeier WE, Xie F, Kurz SG, Lu N, Wood JR. Progressive obesity alters the steroidogenic response to ovulatory stimulation and increases the abundance of mRNAs stored in the ovulated oocyte. *Mol Reprod Dev*. 2014;81(8):735-47.
 10. Mirabi P, Chaichi MJ, Esmailzadeh S, Jorsaraei SG, Bijani A, Ehsani M, et al. The role of fatty acids on ICSI outcomes: a prospective cohort study. *Lipids Health Dis*. 2017;16(1):18.
 11. Haggarty P, Wood M, Ferguson E, Hoar G, Srikantharajah A, Milne E, et al. Fatty acid metabolism in human preimplantation embryos. *Hum Reprod*. 2006;21(3):766-73.
 12. Marei WF, Wathes DC, Fouladi-Nashta AA. Impact of linoleic acid on bovine oocyte maturation and embryo development. *Reproduction*. 2010;139(6):979-88.
 13. McKeegan PJ, Sturmey RG. The role of fatty acids in oocyte and early embryo development. *Reprod Fertil Dev*. 2011;24(1):59-67.
 14. Renaville B, Bacciu N, Comin A, Motta M, Poli I, Vanini G, et al. Plasma and follicular fluid fatty acid profiles in dairy cows. *Reprod Domest Anim*. 2010;45(1):118-21.
 15. Fayezi S, Leroy JL, Novin MG, Darabi M. Oleic acid in the modulation of oocyte and preimplantation embryo development. *Zygote*. 2018;26(1):1-13.
 16. Aardema H, Vos PL, Lolicato F, Roelen BA, Knijn HM, Vaandrager AB, et al. Oleic acid prevents detrimental effects of saturated fatty acids on bovine oocyte developmental competence. *Biol Reprod*. 2011;85(1):62-9.
 17. Jorritsma R, Cesar ML, Hermans JT, Kruitwagen CL, Vos PL, Kruip TA. Effects of non-esterified fatty acids on bovine granulosa cells and developmental potential of oocytes in vitro. *Anim Reprod Sci*. 2004;81(3-4):225-35.
 18. Mennitti LV, Oliveira JL, Morais CA, Estadella D, Oyama LM, do Nascimento CM, et al. Type of fatty acids in maternal diets during pregnancy and/or lactation and metabolic consequences of the offspring. *J Nutr Biochem*. 2015;26(2):99-111.
 19. Dumesic DA, Meldrum DR, Katz-Jaffe MG, Krisner RL, Schoolcraft WB. Oocyte environment: follicular fluid and cumulus cells are critical for oocyte health. *Fertil Steril*. 2015;103(2):303-16.
 20. Mirmiran P, Esfahani FH, Mehrabi Y, Hedayati M, Azizi F. Reliability and relative validity of an FFQ for nutrients in the Tehran lipid and glucose study. *Public Health Nutr*. 2010;13(5):654-62.
 21. Hosseini Esfahani F, Asghari G, Mirmiran P, Azizi F. Reproducibility and relative validity of food group intake in a food frequency questionnaire developed for the Tehran Lipid and Glucose Study. *J Epidemiol*. 2010;20(2):150-8.
 22. Esmailzadeh A, Mirmiran PA, Azizi F. Whole-grain consumption and the metabolic syndrome: a favorable association in Tehranian adults. *Eur J Clin Nutr*. 2005;59(3):353-62.
 23. Craig CL, Marshall AL, Sjöström M, Bauman AE, Booth ML, Ainsworth BE, et al. International physical activity questionnaire: 12-country reliability and validity. *Med Sci Sports Exerc*. 2003;35(8):1381-95.
 24. Moghaddam MB, Aghdam FB, Jafarabadi MA, Allahverdipour H, Nikookheslat SD, Safarpour S. The Iranian version of international physical activity questionnaire (IPAQ) in Iran: content and construct validity, factor structure, internal consistency and stability. *World Appl Sci J*. 2012;18(8):1073-80.
 25. Azadi-Yazdi M, Karimi-Zarchi M, Salehi-Abargouei A, Fallahzadeh H, Nadjarzadeh A. Effects of dietary approach to stop hypertension diet on androgens, antioxidant status and body composition in overweight and obese women with polycystic ovary syndrome: a randomised controlled trial. *J Hum Nutr Diet*. 2017;30(3):275-83.
 26. Langbein A, Bartholomeus E, Leroy JL, Bols PE. Bovine in vitro reproduction models can contribute to the development of (female) fertility preservation strategies. *Theriogenology*. 2015;84(4):477-89.
 27. Valckx SD, Leroy JL. The effect of maternal metabolic health and diet on the follicular fluid composition and potential consequences for oocyte and embryo quality. In: Watson R, editor. *Handbook of Fertility*. USA: Academic Press; 2015. p. 35-44.
 28. Salehi R, Colazo MG, Oba M, Ambrose DJ. A prepartum diet supplemented with oilseeds high in oleic or linoleic acid reduced GnRH-induced LH release in dairy cows during second week postpartum. *Reprod Biol Endocrinol*. 2015;13:69.
 29. Bilby TR, Block J, Do Amaral BC, Sa Filho O, Silvestre FT, Hansen PJ, et al. Effects of dietary unsaturated fatty acids on oocyte quality and follicular development in lactating dairy cows in summer. *J Dairy Sci*. 2006;89(10):3891-903.

30. Roura M, Catalá MG, Soto-Heras S, Hammami S, Izquierdo D, Fouladi-Nashta A, et al. Linoleic (LA) and linolenic (ALA) acid concentrations in follicular fluid of prepubertal goats and their effect on oocyte in vitro maturation and embryo development. *Reprod Fertil Dev.* 2018;30(2):286-96.
31. Pratt HP, George MA. Organisation and assembly of the surface membrane during early cleavage of the mouse embryo. *Roux's Arch Dev Biol.* 1989;198(3):170-8.
32. Veshkini A, Khadem AA, Mohammadi-Sangcheshmeh A, Alamouti AA, Soleimani M, Gastal EL. Linolenic acid improves oocyte developmental competence and decreases apoptosis of in vitro-produced blastocysts in goat. *Zygote.* 2016;24(4):537-48.
33. Lee JE, Yong H, Kim HY, Lee WH, Cheong HT, Yang BK, et al. Effect of alpha-linolenic acid on oocyte maturation and embryo development in pigs. *Dev Reprod.* 2017;21(2):205-13.
34. Marei WF, Wathes DC, Fouladi-Nashta AA. The effect of linolenic acid on bovine oocyte maturation and development. *Biol Reprod.* 2009;81(6):1064-72.
35. Calder MD, Caveney AN, Westhusin ME, Watson AJ. Cyclooxygenase-2 and prostaglandin E2 (PGE2) receptor messenger RNAs are affected by bovine oocyte maturation time and cumulus-oocyte complex quality, and PGE2 induces moderate expansion of the bovine cumulus in vitro. *Biol Reprod.* 2001;65(1):135-40.
36. Abayasekara DR, Wathes DC. Effects of altering dietary fatty acid composition on prostaglandin synthesis and fertility. *Prostaglandins Leukot Essent Fatty Acids.* 1999;61(5):275-87.
37. Murakami K, Chan SY, Routtenberg A. Protein kinase C activation by cis-fatty acid in the absence of Ca²⁺ and phospholipids. *J Biol Chem.* 1986;261(33):15424-9.
38. Frey MR, Clark JA, Leontieva O, Uronis JM, Black AR, Black JD. Protein kinase C signaling mediates a program of cell cycle withdrawal in the intestinal epithelium. *J Cell Biol.* 2000;151(4):763-78.
39. Ciepiela P, Bączkowski T, Drozd A, Kazienko A, Stachowska E, Kurzawa R. Arachidonic and linoleic acid derivatives impact oocyte ICSI fertilization—a prospective analysis of follicular fluid and a matched oocyte in a 'one follicle—one retrieved oocyte—one resulting embryo' investigational setting. *PLoS One.* 2015;10(3):e0119087.
40. Chen S, Yao L, Cunningham TJ. Secreted phospholipase A2 involvement in neurodegeneration: differential testing of pro-survival and anti-inflammatory effects of enzyme inhibition. *PLoS One.* 2012;7(6):e39257.
41. Igosheva N, Abramov AY, Poston L, Eckert JJ, Fleming TP, Duchon MR, et al. Maternal diet-induced obesity alters mitochondrial activity and redox status in mouse oocytes and zygotes. *PLoS One.* 2010;5(4):e10074.
42. Wu LL, Russell DL, Norman RJ, Robker RL. Endoplasmic reticulum (ER) stress in cumulus-oocyte complexes impairs pentraxin-3 secretion, mitochondrial membrane potential ($\Delta\Psi$ m), and embryo development. *Mol Endocrinol.* 2012;26(4):562-73.
43. Wu LL, Dunning KR, Yang X, Russell DL, Lane M, Norman RJ, et al. High-fat diet causes lipotoxicity responses in cumulus-oocyte complexes and decreased fertilization rates. *Endocrinology.* 2010;151(11):5438-45.
44. Leroy JL, Vanholder T, Mateusen B, Christophe A, Opsomer G, de Kruif A, et al. Non-esterified fatty acids in follicular fluid of dairy cows and their effect on developmental capacity of bovine oocytes in vitro. *Reproduction.* 2005;130(4):485-95.
45. Tripathi SK, Nandi S, Gupta PS, Mondal S. Influence of common saturated and unsaturated fatty acids on development of ovine oocytes in vitro. *Asian J Anim Sci.* 2015;9:420-6.
46. Alves JPM, Bertolini M, Bertolini LR, Silva CMG, Rondina D. Lipotoxicity: impact on oocyte quality and reproductive efficiency in mammals. *Anim Reprod Belo Horiz.* 2015;12(2):291-7.
47. Hammiche F, Vujkovic M, Wijburg W, de Vries JH, Macklon NS, Laven JS, et al. Increased pre-conception omega-3 polyunsaturated fatty acid intake improves embryo morphology. *Fertil Steril.* 2011;95(5):1820-3.
48. Broughton KS, Rule DC, Ye Y, Zhang X, Driscoll M, Culver B. Dietary omega-3 fatty acids differentially influence ova release and ovarian cyclooxygenase-1 and cyclooxygenase-2 expression in rats. *Nutr Res.* 2009;29(3):197-205.
49. Mattos R, Staples CR, Williams J, Amorocho A, McGuire MA, Thatcher WW. Uterine, ovarian, and production responses of lactating dairy cows to increasing dietary concentrations of menhaden fish meal. *J Dairy Sci.* 2002;85(4):755-64.
50. Amaya-Montoya C, Matsui M, Kawashima C, Hayashi KG, Matsuda G, Kaneko E, et al. Induction of ovulation with GnRH and PGF(2 alpha) at two different stages during the early postpartum period in dairy cows: ovarian response and changes in hormone concentrations. *J Reprod Dev.* 2007;53(4):867-75.
51. Wathes DC, Cheng Z, Mareiy W, Fouladi-Nashta

- A. Polyunsaturated fatty acids and fertility in female mammals: An update. *CAB Rev Perspect Agric Vet Sci Nutr Nat Resour*. 2013;8(041):1–14.
52. Shahnazi V, Zaree M, Nouri M, Mehrzad-Sadaghiani M, Fayezi S, Darabi M, et al. Influence of ω -3 fatty acid eicosapentaenoic acid on IGF-1 and COX-2 gene expression in granulosa cells of PCOS women. *Iran J Reprod Med*. 2015;13(2):71-8.
53. Toori MA, Mosavi E, Nikseresht M, Barmak MJ, Mahmoudi R. Influence of insulin-like growth factor-i on maturation and fertilization rate of immature oocyte and embryo development in NMRI mouse with TCM199 and α -MEM medium. *J Clin Diagn Res*. 2014;8(12):AC05-8.
54. Cerri RL, Juchem SO, Chebel RC, Rutigliano HM, Bruno RG, Galvao KN, et al. Effect of fat source differing in fatty acid profile on metabolic parameters, fertilization, and embryo quality in high-producing dairy cows. *J Dairy Sci*. 2009;92(4):1520-31.
55. Thangavelu G, Colazo MG, Ambrose DJ, Oba M, Okine EK, Dyck MK. Diets enriched in unsaturated fatty acids enhance early embryonic development in lactating Holstein cows. *Theriogenology*. 2007;68(7):949-57.
56. Gueyffier F, Cucherat M. The limitations of observation studies for decision making regarding drugs efficacy and safety. *Therapie*. 2019;74(2): 181-5.